



MVP Health Care Medical Policy

Medicare Part B: Zulresso™ (brexanolone)

Type of Policy:	Medical Therapy
Prior Approval Date:	N/A
Approval Date:	11/01/2023
Effective Date:	01/01/2023
Related Policies:	NA

Codes Requiring Prior Authorization (covered under the medical benefit)

J1632 Zulresso™ (brexanolone)

Overview/Summary of Evidence

Peripartum Depression (formerly Postpartum depression (PPD)) is a mood disorder that can occur during pregnancy or after childbirth which is accompanied by persistent, intense feelings of anxiety, despair or sadness and changes in energy, sleep, and appetite. The symptoms can interfere with a mother's daily tasks and taking care of their child(ren). PPD is different from "baby blues", which is a common occurrence that subsides within a few days to 1-2 weeks without treatment. An estimated one in seven women experience peripartum depression. Current treatment includes psychotherapy (counseling or "talk therapy"), antidepressants or both. Zulresso™ (brexanolone) is the first available medication specifically indicated for the treatment of postpartum depression (PPD) in adults. It is a neuroactive steroid gamma-aminobutyric acid (GABA) A receptor positive modulator that is administered as a one-time continuous intravenous infusion given over 60 hours (2.5 days). Zulresso must be administered in an inpatient setting as it requires continuous monitoring by a healthcare provider for hypoxia, loss of consciousness and excessive sedation. Zulresso is only available through a REMS program.

Indications/Criteria

Zulresso will be considered for coverage when all the following are met:

- Must be 15 years old or older
- Must be female
- Member is ≤ 12 months postpartum
- Must have a confirmed diagnosis of PPD
 - Documentation of baseline Hamilton Depression Rating Scale (HAM-D) score ≥ 20
 - Chart notes documenting a diagnosis of PPD

- Symptom onset within third trimester of pregnancy or within 4 weeks of delivery
- Documentation of a failure, contraindication, or intolerance to at least a 4-week trial of first line antidepressant therapy (i.e. SSRI (sertraline, escitalopram), mirtazapine, venlafaxine, duloxetine) at the maximally tolerated FDA-approved dose **OR**
- If a 4-week trial with an oral antidepressant is inappropriate, clinical rationale must be documented in the medical record and will be considered on a case by case basis (such as cases of severe PPD) Limit one treatment course per postpartum period- one continuous 60-hour intravenous infusion with a healthcare provider available on site to continuously monitor the patient at a healthcare setting that is certified in the REMS program.

Approval will be for one infusion per postpartum period

Exclusions

- End stage renal disease (ESRD)
 - Home infusion
 - Male
 - Concurrent active psychosis, bipolar disorder and schizoaffective disorder
 - Multiple infusions in the same postpartum period
 - Pregnant
-

References

1. Zulresso TM (brexanolone) injection, for intravenous use. Prescribing Information. Cambridge, MA. Sage Therapeutics. June 2022.
2. American College of Obstetricians and Gynecologists. Frequently Asked Question Labor, Delivery, and Postpartum Care. December 2013. Available at: <https://www.acog.org/Patients/FAQs/Postpartum-Depression>
3. National Institute of Mental Health. Postpartum Depression Facts. Available at: <https://www.nimh.nih.gov/health/publications/postpartum-depression-facts/index.shtml#pub2>
4. Melzer-Brody S, Colquhoun H, Reisenberg R, et al. Brenaxolone injection in postpartum depression: two multicenter, double-blind, randomized, placebo-controlled, phase 3 trials. *Lancet*. 2018; 392(10152):1058-1070.
5. Frieder A, Fersh M, Hainline R, et al. Pharmacotherapy of Postpartum Depression: Current Approaches and Novel Drug Development. *CNS Drugs*. 2019 Feb; 33(1): 265-282.

6. Williams J, Ryan D, Thomas-Peter K, et al. Best Practice Guidelines for Mental Health Disorders in the Perinatal Period. BC Reproductive Mental Health Program & Perinatal Services. 2014 Mar; 13-89
7. Yonkers KA, Wisner KL, Stewart DE, et al. The management of depression during pregnancy: a report from the American Psychiatry Association and the American College of Obstetricians and Gynecologists. *Gen Hosp Psychiatry* 2009;31:403-13

Davé S1, Petersen I, Sherr L, Nazareth I. Incidence of maternal and paternal depression in primary care: a cohort study using a primary care database. *Arch Pediatr Adolesc Med*. 2010 Nov;164(11):1038-44.