



# 2019 Coding Reference Guide for Primary Care



This HEDIS Coding document is updated annually.  
Please visit [ncqa.org/hedis/measures](https://www.ncqa.org/hedis/measures) for updates throughout 2019.

## 2019 HEDIS Coding Reference Guide Contents

### Antibiotic Measures ..... 1

Appropriate Testing for Children with Pharyngitis (CWP) (Admin*) .....	1
Appropriate Treatment for Children with Upper Respiratory Infection (URI) (Admin) .....	1
Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis (AAB) (Admin) .....	1

### Preventive Care Measures ..... 3

Adolescent Well-Care Visits (AWC) (Admin) .....	3
Childhood Immunization Status (CIS).....	3
Lead Screening in Children (LSC) .....	4
Immunizations for Adolescents (IMA) (Hybrid**) .....	4
Adolescent Preventive Care (ADL) (Hybrid) QARR .....	4
Adult BMI Assessment (ABA) (Hybrid) .....	5
Well-Child Visits in the First 15 Months of Life (W15) (Admin) .....	5
Well-Child Visits in the Third, Fourth, Fifth, and Sixth Years of Life (W34) .....	5
Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents (WCC) (Hybrid).....	6
Breast Cancer Screening (BCS) (Admin) .....	6
Chlamydia Screening in Women (CHL) (Admin) .....	6
Cervical Cancer Screening (CCS) (Hybrid).....	6
Colorectal Cancer Screening (COL) (Hybrid) .....	6

### Testing/Screening Measures ..... 7

Use of Spirometry Testing in the Assessment and Diagnosis of COPD (SPR) (Admin) .....	7
Use of Imaging Studies for Low Back Pain (LBP) (Admin).....	7

### Disease Management Measures..... 8

Medication Management for People with Asthma (MMA) (Admin) .....	8
Asthma Medication Ratio (AMR) (Admin) .....	8
Controlling High Blood Pressure (CBP) (Hybrid).....	8
Osteoporosis Management in Women Who Had a Fracture (OMW) (Admin).....	9
Medication Reconciliation Post-Discharge (MRP) (Admin).....	10
Disease-Modifying Anti-Rheumatic Drug Therapy for Rheumatoid Arthritis (ART) (Admin) .....	10
Pharmacotherapy Management of COPD Exacerbation (PCE) (Admin) .....	11
Comprehensive Diabetes Care (CDC) (Hybrid).....	12
Medical Attention for Nephropathy—Urine Protein Test.....	13
Medical Attention for Nephropathy—Nephropathy Treatment.....	13

### Medication Management Measures..... 14

Annual Monitoring for Patients on Persistent Medications (MPM) (Admin).....	14
Persistence of Beta-Blocker Treatment After a Heart Attack (PBH) (Admin).....	15

\* Administrative (Admin) measures are calculated using claims data only.

\*\* Hybrid measures are calculated using both claims data and data gathered from medical record review.

Current Procedural Terminology ©2014 American Medical Association. All rights reserved. Current Procedural Terminology (CPT) and Uniform Bill (UB) codes are stated to the minimum specificity required.

Appropriate Care and Associated Codes to Bill			
	CPT and CPT II	ICD-10	Goal
<p><b>Appropriate Testing for Children with Pharyngitis (CWP) (Admin*)</b></p> <p>The percentage of children 3–18 years of age who were diagnosed with pharyngitis, dispensed an antibiotic, and received a group A streptococcus (strep) test for the episode. A higher rate represents better performance (i.e., appropriate testing).</p>	<p><b>Acceptable Strep Test Codes</b> 87070, 87071, 87081, 87430, 87650, 87651, 87652, 87880 <i>(MVP Medicaid Members only)</i></p>	<p><b>Streptococcal Pharyngitis</b> J02.0</p> <p><b>Acute Pharyngitis Due to Other Specified Organisms</b> J02.8</p> <p><b>Acute Pharyngitis, Unspecified</b> J02.9</p> <p><b>Acute Streptococcal Tonsillitis, Unspecified</b> J03.00</p> <p><b>Acute Recurrent Streptococcal Tonsillitis</b> J03.01</p> <p><b>Acute Tonsillitis Due to Other Specified Organisms</b> J03.80</p> <p><b>Acute Recurrent Tonsillitis Due to Other Specified Organisms</b> J03.81</p> <p><b>Acute Tonsillitis, Unspecified</b> J03.90</p> <p><b>Acute Recurrent Tonsillitis, Unspecified</b> J03.91</p>	<p>To perform a group A streptococcus test and provide counseling to patients regarding the inappropriate use of antibiotics.</p>
<p><b>Appropriate Treatment for Children with Upper Respiratory Infection (URI) (Admin)</b></p> <p>The percentage of children 3 months–18 years of age who were given a diagnosis of upper respiratory infection (URI) and were not dispensed an antibiotic prescription.</p>		<p><b>Acute Nasopharyngitis (Common Cold)</b> J00</p> <p><b>Acute Laryngopharyngitis</b> J06.0</p> <p><b>Acute Upper Respiratory Infection, Unspecified</b> J06.9</p>	<p>Provide counseling to patients regarding the inappropriate use of antibiotics.</p>
<p><b>Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis (AAB) (Admin)</b></p> <p>The percentage of adults 18–64 years of age with a diagnosis of acute bronchitis who were not dispensed an antibiotic prescription.</p>		<p><b>Acute Bronchitis Due to Coxsackievirus</b> J20.3</p> <p><b>Acute Bronchitis Due to Parainfluenza Virus</b> J20.4</p> <p><b>Acute Bronchitis Due to Hemophilus Influenza</b> J20.1</p> <p><b>Acute bronchitis Due to Streptococcus</b> J20.2</p> <p><b>Bronchitis, Not Specified as Acute or Chronic</b> J40</p> <p><i>(Continued on page 2)</i></p>	<p>Provide counseling to patients regarding the inappropriate use of antibiotics and to not prescribe an antibiotic without indication of a competing diagnosis or co-morbidity.</p>

\*Administrative (Admin) measures are calculated using claims data only.

Appropriate Care and Associated Codes to Bill		
CPT and CPT II	ICD-10	Goal
<p><i>Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis (AAB) (Admin) continued from page 1</i></p>	<p><i>(Continued from page 1)</i></p> <p><b>Acute Laryngopharyngitis</b> J06.0</p> <p><b>Acute Upper Respiratory Infection, Unspecified</b> J06.9</p> <p><b>Acute Bronchitis Due to Mycoplasma Pneumoniae</b> J20.0</p> <p><b>Acute Bronchitis Due to Respiratory Syncytial Virus</b> J20.5</p> <p><b>Acute Bronchitis Due to Rhinovirus</b> J20.6</p> <p><b>Acute Bronchitis Due to Echovirus</b> J20.7</p> <p><b>Acute Bronchitis Due to Other Specified Organisms</b> J20.8</p> <p><b>Acute Bronchitis, Unspecified</b> J20.9</p>	

	Appropriate Care and Associated Codes to Bill			
	CPT and CPT II	ICD-10	HCPCS	UB Revenue
<p><b>Adolescent Well-Care Visits (AWC) (Admin)</b></p> <p>The percentage of enrolled members 12–21 years of age who had at least one comprehensive well-care visit with a PCP or an OB/GYN practitioner during the measurement year.</p> <p><i>Use either CPT, ICD-10, or HCPCS codes.</i></p>	<p>99381, 99382, 99383, 99384, 99385, 99391, 99392, 99393, 99394, 99395, 99461</p>	<p>Z00.00, Z00.01, Z00.110, Z00.111, Z00.121, Z00.129, Z00.5, Z00.8, Z02.0, Z02.1, Z02.2, Z02.3, Z02.4, Z02.5, Z02.6, Z02.71, Z02.82</p> <p><b>Encounter for Health Supervision and Care of Foundling</b> Z76.1</p> <p><b>Encounter for Health Supervision and Care of Other Healthy Infant and Child</b> Z76.2</p>	<p>G0438, G0439</p>	
<p><b>Childhood Immunization Status (CIS)</b></p> <p>The percentage of children two years of age who had all of the following vaccines by their second birthday:</p> <ul style="list-style-type: none"> <li>• four diphtheria, tetanus, and acellular pertussis (DTaP)</li> <li>• three polio (IPV)</li> <li>• one measles, mumps, and rubella (MMR)</li> <li>• three haemophilus influenza type B (HiB)</li> <li>• three hepatitis B (HepB)</li> <li>• one chicken pox (VZV)</li> <li>• four pneumococcal conjugate (Pcv)</li> <li>• one hepatitis A (Hep A)</li> <li>• two or three rotavirus (RV)</li> <li>• two influenza (flu)</li> </ul> <p>The measure calculates a rate for each vaccine and nine separate combination rates (Hybrid).</p>	<p><b>DTaP</b> 90698, 90700, 90721, 90723</p> <p><b>HiB</b> 90644, 90645, 90646, 90647, 90648, 90698, 90721, 90748</p> <p><b>Hep A</b> 90633</p> <p><b>Hep B</b> 90723, 90740, 90744, 90747, 90748</p> <p><b>IPV</b> 90698, 90713, 90723</p> <p><b>Influenza</b> 90655, 90657, 90661, 90662, 90673, 90685, 90686, 90687, 90688</p> <p><b>Measles</b> 90705</p> <p><b>Mumps</b> 90704</p> <p><b>Rubella</b> 90706</p> <p><b>Measles/Rubella</b> 90708</p> <p><b>MMR</b> 90707, 90710</p> <p><b>PCV</b> 90669, 90670</p> <p><b>Rotavirus (two dose)</b> 90681</p> <p><b>Rotavirus (three dose)</b> 90680</p> <p><b>VZV</b> 90710, 90716</p>		<p><b>Hep B</b> G0010</p> <p><b>Influenza</b> G0008</p> <p><b>PCV</b> G0009</p>	

	Appropriate Care and Associated Codes to Bill			
	CPT and CPT II	ICD-10	HPCPS	UB Revenue
<p><b>Lead Screening in Children (LCS)</b></p> <p>The percentage of children two years of age who had one or more capillary or venous lead blood test for lead poisoning by their second birthday.</p>	83655			
<p><b>Immunizations for Adolescents (IMA) (Hybrid**)</b></p> <p>The percentage of adolescents 13 years of age who had all of the following vaccines by their 13th birthday:</p> <ul style="list-style-type: none"> <li>one dose of meningococcal conjugate vaccine</li> <li>one tetanus, diphtheria toxoids, and acellular pertussis (Tdap) vaccine</li> <li>completed the human papillomavirus (HPV)</li> </ul> <p>The measure calculates a rate for each vaccine and two combination rates.</p>	<p><b>Meningococcal</b> 90734</p> <p><b>Tdap</b> 90715</p> <p><b>HPV</b> 90649, 90650, 90651</p>			
<p><b>Adolescent Preventive Care (ADL) (Hybrid) QARR</b></p> <p>The percentage of members 12–17 years of age who had at least one visit with the PCP or OB/GYN during the measurement year, and received the following four components of care during the measurement year:</p> <ol style="list-style-type: none"> <li>Assessment, counseling, or education on risk behaviors and preventive actions associated with sexual activity during the measurement year. Risk behaviors and preventative actions for sexual activity include: abstinence, current behaviors, family planning, condom use, contraceptives, HIV, STIs, pregnancy prevention, and safe sex.</li> <li>Assessment, counseling, or education about depression during the measurement year. Depression has an affective component (mood, interest, and enjoyment) and a physical component (changes in appetite, sleep pattern, and concentration). Use of assessment tool or provider interview have been determined to be more effective methods for identification of depression than relying on patient self-report.</li> <li>Assessment, counseling, or education about the risks of tobacco use during the measurement year. Tobacco includes, but is not limited to, cigarettes, cigars, chew, or other forms of smokeless tobacco.</li> <li>Assessment, counseling, or education about the risks of substance use during the measurement year. Substance use includes, but is not limited to, alcohol, street drugs, non-prescription drugs, prescription drugs misuse, and inhalant use.</li> </ol>	<p><b>Screening for High Risk Sexual Behavior</b> 4293F</p> <p><b>Depression</b> 1220F, 3085F, 3351F, 3352F, 3353F, 3354F, 3725F</p> <p><b>Tobacco Use Assessment</b> 1000F, 1031F, 1032F, 1033F, 1034F, 1035F, 1036F, 99406, 99407</p> <p><b>Tobacco Cessation Counseling or Services</b> 99406, 99407, 4000F, 4001F, 4004F</p> <p><b>Alcohol and/or Drug Assessment or Screening</b> 99408, 99409, 3016F, 4290F</p> <p><b>Alcohol and/or Drug Use Counseling Services</b> 4306F, 4320F</p>	<p><b>Counseling for HIV</b> Z71.7</p> <p><b>Counseling for Other STIs</b> Z70.8</p> <p><b>Counseling for Oral and Other Contraceptives</b> Z30.0, Z30.01, Z30.011, Z30.012, Z30.013, Z30.014, Z30.015, Z30.016, Z30.017, Z30.018, Z30.019, Z30.02, Z30.09</p> <p><b>Screening for High Risk Sexual Behavior</b> Z72.5, Z72.51, Z72.53, Z72.53, Z70, Z70.1, Z70.2, Z70.3, Z70.9</p> <p><b>Tobacco Cessation Counseling or Services</b> Z71.6</p> <p><b>Alcohol and/or Drug Use Counseling Services</b> Z71.41, Z71.51</p>	<p><b>Screening for High Risk Sexual Behavior</b> G0445</p> <p><b>Depression</b> G0444, G8431, G8510, G8511, G8930 S3005</p> <p><b>Tobacco Cessation or Counseling Services</b> G0436, G0437</p> <p><b>Tobacco Cessation Classes</b> S9453</p> <p><b>Alcohol and/or Drug Assessment or Screening</b> G0396, G0397, H0001, H0049</p> <p><b>Alcohol and/or Drug Use Counseling Services</b> G0443 H0005, H0006, H0007, H0022, H0047, H0050, T1007</p>	

\*\*Hybrid measures are calculated using both claims data and data gathered from medical record review.

	Appropriate Care and Associated Codes to Bill			
	CPT and CPT II	ICD-10	HCPCS	UB Revenue
<p><b>Adult BMI Assessment (ABA) (Hybrid)</b></p> <p>The percentage of members 18–74 years of age who had an outpatient visit and whose body mass index (BMI) was documented during the measurement year or the year prior to the measurement year.</p> <p>20 years of age or older: BMI percentile and weight 19 years of age or younger: BMI percentile, weight, and height</p>	<p>99201, 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99241, 99242, 99243, 99244, 99245, 99341, 99342, 99343, 99344, 99345, 99347, 99348, 99349, 99350, 99381, 99382, 99383, 99384, 99385, 99386, 99387, 99391, 99392, 99393, 99394, 99395, 99396, 99397, 99401, 99402, 99403, 99404, 99411, 99412, 99429, 99455, 99456</p>	<p><b>BMI</b> Z68.1, Z68.20, Z68.21, Z68.22, Z68.23, Z68.24, Z68.25, Z68.26, Z68.27, Z68.28, Z68.29, Z68.30, Z68.31, Z68.32, Z68.33, Z68.34, Z68.35, Z68.36, Z68.37, Z68.38, Z68.39, Z68.41, Z68.42, Z68.43, Z68.44, Z68.45</p> <p><b>BMI Percentile</b> Z68.51, Z68.52, Z68.53, Z68.54</p>	<p><b>Initial Preventive Physical Exam Visit for New Beneficiary During the First 12 Months of Medicare Enrollment</b></p> <p><b>Annual Wellness Visit—Includes a Personalized Prevention Plan of Service (PPS), Initial Visit</b> G0438</p> <p><b>Annual Wellness Visit—Includes a Personalized Prevention Plan of Service (PPS), Subsequent Visit</b> G0439</p> <p><b>Hospital Outpatient Clinic Visit for Assessment and Management of a Patient</b> G0463</p> <p><b>Clinic Visit/Encounter, All-Inclusive</b> T1015</p>	<p>0510, 0511, 0512, 0513, 0514, 0515, 0516, 0517, 0519, 0520, 0521, 0522, 0523, 0526, 0527, 0528, 0529, 0982, 0983</p>
<p><b>Well-Child Visits in the First 15 Months of Life (W15) Admin</b></p> <p>The percentage of members who turned 15 months old during the measurement year and who had six or more well-child visits with a PCP during their first 15 months of life.</p> <p><i>Use either CPT, ICD-10, or HCPCS codes.</i></p>	<p>99381, 99382, 99383, 99384, 99385, 99391, 99392, 99393, 99394, 99395, 99461</p>	<p>Z00.00, Z00.01, Z00.110, Z00.111, Z00.121, Z00.129, Z00.5, Z00.8, Z02.0, Z02.1, Z02.2, Z02.3, Z02.4, Z02.5, Z02.6, Z02.71, Z02.82</p> <p><b>Encounter for Health Supervision and Care of Foundling</b> Z76.1</p> <p><b>Encounter for Health Supervision and Care of Other Healthy Infant and Child</b> Z76.2</p>	<p>G0438, G0439</p>	
<p><b>Well-Child Visits in the Third, Fourth, Fifth, and Sixth Years of Life (W34) Admin</b></p> <p>The percentage of members 3–6 years of age who had one or more well-child visits with a PCP during the measurement year.</p> <p><i>Use either CPT, ICD-10, or HCPCS codes.</i></p>	<p>99381, 99382, 99383, 99384, 99385, 99391, 99392, 99393, 99394, 99395, 99461</p>	<p>Z00.00, Z00.01, Z00.110, Z00.111, Z00.121, Z00.129, Z00.5, Z00.8, Z02.0, Z02.1, Z02.2, Z02.3, Z02.4, Z02.5, Z02.6, Z02.71, Z02.82</p> <p><b>Encounter for Health Supervision and Care of Foundling</b> Z76.1</p> <p><b>Encounter for Health Supervision and Care of Other Healthy Infant and Child</b> Z76.2</p>	<p>G0438, G0439</p>	



	Appropriate Care and Associated Codes to Bill			
	CPT and CPT II	ICD-10	HCPCS	UB Revenue
<p><b>Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents (WCC) (Hybrid)</b></p> <p>The percentage of members 3–17 years of age who had an outpatient visit with a PCP or OB/GYN, and who had evidence of the following during the measurement year:</p> <ul style="list-style-type: none"> <li>• BMI Percentile Documentation</li> <li>• Counseling for Nutrition</li> <li>• Counseling for Physical Activity</li> </ul>	<p><b>Nutrition Counseling</b> 97802, 97803, 97804</p>	<p><b>BMI Percentile</b> Z68.51 BMI &lt;5th percentile Z68.52 BMI 5th–85th percentile Z68.53 BMI 85th–95th percentile Z68.54 BMI &gt;95th percentile Z71.3 (Dietary Surveillance and Counseling)</p> <p><b>Physical Activity</b> Z02.5</p> <p><b>Exercise Counseling</b> Z71.82</p>	<p><b>Nutrition Counseling</b> G0270, G0271, G0447 S9449, S9452, S9470</p> <p><b>Physical Activity</b> G0447 S9451</p>	
<p><b>Breast Cancer Screening (BCS) (Admin)</b></p> <p>The percentage of women 50–74 years of age who had a mammogram to screen for breast cancer any time on or between October 1, 2016 and December 31, 2018.</p>	<p>77055, 77056, 77057, 77061, 77062, 77063, 77065, 77066, 77067</p>		<p>G0202, G0204, G0206</p>	<p>0401, 0403</p>
<p><b>Chlamydia Screening in Women (CHL) (Admin)</b></p> <p>The percentage of women 16–24 years of age who were identified as sexually active and who had at least one test for Chlamydia during the measurement year.</p>	<p><b>Chlamydia Tests</b> 87110, 87270, 87320, 87490, 87491, 87492, 87810</p>			
<p><b>Cervical Cancer Screening (CCS) (Hybrid)</b></p> <p>The percentage of women 21–64 years of age who were screened for cervical cancer by <i>either</i> of the following:</p> <ul style="list-style-type: none"> <li>• Women age 21–64 who had cervical cytology performed every 3 years.</li> <li>• Women age 30–64 who had cervical cytology/human Papillomavirus (HPV) co-testing performed every 5 years</li> </ul>	<p><b>Cervical Cytology</b> 88141, 88142, 88143, 88147, 88148, 88150, 88152, 88153, 88154, 88164, 88165, 88166, 88167, 88174, 88175</p> <p><b>HPV Test</b> 87620, 87621, 87622, 87624, 87625</p>		<p><b>Cervical Cytology</b> G0123, G0124, G0141, G0143, G0144, G0145, G0147, G0148 P3000, P3001</p> <p><b>HPV Test</b> G0476</p>	<p><b>Cervical Cytology</b> 0923</p>
<p><b>Colorectal Cancer Screening (COL) (Hybrid)</b></p> <p>The percentage of members 50–75 years of age who had one or more colorectal cancer screenings:</p> <ul style="list-style-type: none"> <li>• Colonoscopy during the period 2009–2018</li> <li>• CT-Colonography during the period 2014–2018</li> <li>• FIT-DNA during the period 2016–2018</li> <li>• Flexible sigmoidoscopy during the period 2014–2018</li> <li>• FOBT during 2018</li> </ul>	<p><b>Fecal Occult Blood Test (FOBT)</b> 82270, 82274</p> <p><b>FIT-DNA</b> 81528</p> <p><b>Flexible Sigmoidoscopy</b> 45330, 45331, 45332, 45333, 45334, 45335, 45337, 45338, 45339, 45340, 45341, 45342, 45345, 45346, 45347, 45349, 45350</p> <p><b>Colonoscopy</b> 44388–44394, 44397, 44401–44408, 45355, 45378–45393, 45398</p> <p><b>CT-Colonography</b> 74261, 74262, 74263</p>		<p><b>Fecal Occult Blood Test (FOBT)</b> G0328</p> <p><b>FIT-DNA</b> G0464</p> <p><b>Flexible Sigmoidoscopy</b> G0104</p> <p><b>Colonoscopy</b> G0105, G0121</p>	



	Appropriate Care and Associated Codes to Bill			
	CPT and CPT II	ICD-10	HCPCS	UB Revenue
<p><b>Use of Spirometry Testing in the Assessment and Diagnosis of COPD (SPR) (Admin)</b></p> <p>The percentage of members 40 years of age and older with a new diagnosis or newly active COPD who received appropriate spirometry testing to confirm the diagnosis.</p>	<p><b>Spirometry</b> 94010, 94014, 94105, 94016, 94060, 94070, 94375, 94620</p>	<p><b>Chronic Bronchitis</b> J41.0, J41.1, J41.8, J42</p> <p><b>Emphysema</b> J43.0, J43.1, J43.2, J43.8, J43.9</p> <p><b>COPD (including Dx codes listed above for Chronic Bronchitis and Emphysema)</b> J44.0, J44.1, J44.9, J47.0, J47.1, J47.9</p>		
<p><b>Use of Imaging Studies for Low Back Pain (LBP) (Admin)</b></p> <p>The percentage of members with a primary diagnosis of low back pain that did not have an imaging study (plain X-ray, MRI, CT scan) within 28 days of the diagnosis.</p> <p>A higher score indicates appropriate treatment of low back pain (i.e., the proportion for whom imaging studies did not occur).</p>		<p><b>Uncomplicated Low Back Pain</b> M47.26, M47.27, M47.28, M47.816, M47.817, M47.818, M47.896, M47.897, M47.898, M48.06, M48.061, M48.062, M48.07, M48.08, M51.16, M51.17, M51.26, M51.27, M51.36, M51.37, M51.86, M51.87, M53.2X6, M53.2X7, M53.2X8, M53.3, M53.86, M53.87, M53.88, M54.16, M54.17, M54.18, M54.30, M54.31, M54.32, M54.40, M54.41, M54.42, M54.5, M54.89, M54.9, M99.03, M99.04, M99.23, M99.33, M99.43, M99.53, M99.63, M99.73, M99.83, M99.84 S33.100A, S33.100D, S33.100S, S33.110A, S33.110D, S33.110S, S33.120A, S33.120D, S33.120S, S33.130A, S33.130D, S33.130S, S33.140A, S33.140D, S33.140S, S33.5XXA, S33.6XXA, S33.8XXA, S33.9XXA, S39.002A, S39.002D, S39.002S, S39.012A, S39.012D, S39.012S, S39.092A, S39.092D, S39.092S, S39.82XA, S39.82XD, S39.82XS, S39.92XA, S39.92XD, S39.92XS</p>		

	Appropriate Care and Associated Codes to Bill		
	CPT and CPT II	ICD-10	Prescription
<p><b>Medication Management for People with Asthma (MMA) (Admin)</b></p> <p>The percentage of members 5–64 years of age during the measurement year who were identified as having persistent asthma and were dispensed appropriate medications that they remained on during the treatment period. Two rates are reported:</p> <ol style="list-style-type: none"> <li>1. The percentage of members who remained on an asthma controller medication for at least 50% of the treatment period.</li> <li>2. The percentage of members who remained on an asthma controller medication for at least 75% of the treatment period.</li> </ol>		<p><b>Asthma</b>                      J45.20, J45.21, J45.22, J45.30, J45.31, J45.32, J45.40, J45.41, J45.42, J45.50, J45.51, J45.52, J45.901, J45.902, J45.909, J45.990, J45.991, J45.998</p>	<p><b>Asthma Controller Medications</b></p> <p><b>Antiasthmatic Combinations</b>                      Dyphylline-guaifenesin,                      Guaifenesin-theophylline</p> <p><b>Antibody Inhibitors</b>                      Omalizumab</p> <p><b>Inhaled Steroid Combinations</b>                      Budesonide-formoterol, Fluticasone-salmeterol, Fluticasone-vilanterol, Mometasone-formoterol</p> <p><b>Inhaled Corticosteroids</b>                      Beclomethasone, Budesonide, Ciclesonide, Flunisolide, Fluticasone CFC free, Mometasone</p> <p><b>Leukotriene Modifiers</b>                      Montelukast, Zafirlukast, Zileuton</p> <p><b>Mast Cell Stabilizers</b>                      Cromolyn</p> <p><b>Methylxanthines</b>                      Aminophylline, Dyphylline, Theophylline</p> <p><b>Asthma Reliever Medications</b></p> <p><b>Short-Acting, Inhaled Beta-2 Agonists</b>                      Albuterol, Levalbuterol, Pirbuterol</p>
<p><b>Asthma Medication Ratio (AMR) (Admin)</b></p> <p>The percentage of members 5–64 years of age who were identified as having persistent asthma and had a ratio of controller medications to total asthma medications of 0.50 or greater during the measurement year.</p>		<p>J45.20, J45.21, J45.22, J45.30, J45.31, J45.32, J45.40, J45.41, J45.42, J45.50, J45.51, J45.52, J45.901, J45.902, J45.909, J45.990, J45.991, J45.998</p>	
<p><b>Controlling High Blood Pressure (CBP) (Hybrid)</b></p> <p>The percentage of members 18–85 years of age who had a diagnosis of hypertension (HTN) and whose BP was adequately controlled (&lt;140/90 mm Hg) during the measurement year.</p>		<p><b>Essential (Primary) Hypertension</b>                      I10</p>	

Appropriate Care and Associated Codes to Bill					
	CPT and CPT II	ICD-10	HCPCS	Prescription	UB Revenue
<p><b>Osteoporosis Management in Women Who had a Fracture (OMW) (Admin)</b></p> <p>The percentage of women 67–85 years of age who suffered a fracture and who had either a bone mineral density (BMD) test or prescription for a drug to treat osteoporosis in the six months after the fracture.</p>	<p><b>Fracture</b>                      21811, 21812, 21813, 21820, 21825, 22310, 23500, 23505, 23515, 23570, 23575, 23585, 23600, 23605, 23615, 23616, 23620, 23625, 23630, 24500, 24505, 24515, 24516, 24530, 24535, 24538, 24545, 24546, 24560, 24565, 24566, 24575, 24576, 24577, 24579, 24582, 24650, 24655, 24665, 24666, 24670, 24675, 24685, 25500, 25505, 25515, 25520, 25525, 25526, 25530, 25535, 25545, 25560, 25565, 25574, 25575, 25600, 25605, 25606, 25607, 25608, 25609, 25622, 25624, 25628, 25630, 25635, 25645, 25650, 25651, 25652, 25680, 25685, 26600, 26605, 26607, 26608, 26615, 27200, 27202, 27215, 27220, 27222, 27226, 27227, 27228, 27230, 27232, 27235, 27236, 27238, 27240, 27244, 27245, 27246, 27248, 27254, 27267, 27268, 27269, 27500, 27501, 27502, 27503, 27506, 27507, 27508, 27509, 27510, 27511, 27513, 27514, 27520, 27524, 27530, 27532, 27535, 27536, 27538, 27540, 27750, 27752, 27756, 27758, 27759, 27760, 27762, 27766, 27767, 27768, 27769, 27780, 27781, 27784, 27786, 27788, 27792, 27808, 27810, 27814, 27816, 27818, 27822, 27823, 27824, 27825, 27826, 27827, 27828, 28400, 28405, 28406, 28415, 28420, 28430, 28435, 28436, 28445, 28450, 28455, 28456, 28465, 28470, 28475, 28476, 28485, 29850, 29851, 29855, 29856</p> <p><b>Bone Mineral Density Test</b>                      76977, 77078, 77080, 77081, 77082, 77085, 77086</p> <p><b>Outpatient</b>                      99201, 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99241, 99242, 99243, 99244, 99245, 99341, 99342, 99343, 99344, 99345, 99347, 99348, 99349, 99350, 99381, 99382, 99383, 99384, 99385, 99386, 99387, 99391, 99392, 99393, 99394, 99395, 99396, 99397, 99401, 99402, 99403, 99404, 99411, 99412, 99429, 99455, 99456</p>	<p><b>Fracture</b>                      Code Ranges beginning with BP48ZZ1, BP49ZZ1, BP4GZZ1, BP4LZZ1, BP4MZZ1, BP4NZZ1, BPB4PZZ1, BQ00ZZ1, BQ01ZZ1, BQ03ZZ1, BQ04ZZ1, BR00ZZ1, BR07ZZ1, BR09ZZ1, BR0GZZ1 M48, M80, M84, M97, M180 S12, S22, S32, S42, S49, S52, S59, S62, S72, S79, S82, S89, S92, S99</p>	<p><b>Fracture</b>                      S2360</p> <p><b>Bone Mineral Density Tests</b>                      G0130</p> <p><b>Outpatient</b>                      G0402, G0438, G0439, G0463, T1015</p> <p><b>Injection, Denosumab, 1 mg</b>                      J0897</p> <p><b>Injection, Ibandronate Sodium, 1 mg</b>                      J1740</p> <p><b>Injection, Teriparatide, 10 mcg</b>                      J3110</p> <p><b>Injection, Zoledronic Acid, 1 mg</b>                      J3489</p>	<p><b>Osteoporosis Medications</b></p> <p><b>Biphosphonates</b>                      Alendronate, Alendronate-cholecalciferol, Ibandronate, Risedronate, Zoledronic acid</p> <p><b>Other Agents</b>                      Abaloparatide, Calcitonin, Calcitonin Salmon, Denosumab, Raloxifene, Teriparatide</p>	<p>0510, 0511, 0512, 0513, 0514, 0515, 0516, 0517, 0519, 0520, 0521, 0522, 0523, 0526, 0527, 0528, 0529, 0982, 0983</p>

Appropriate Care and Associated Codes to Bill					
	CPT and CPT II	ICD-10	HCPCS	Prescription	UB Revenue
<p><b>Medication Reconciliation Post-Discharge (MRP) (Admin)</b></p> <p>The percentage of discharges January 1–December 1 of the measurement year for members 18 years of age and older for whom medications were reconciled the date of discharge through 30 days after discharge (31 total days)</p>	<p>Medication Reconciliation CPT: 99483, 99495, 99496 CPT-CAT-II: 1111F</p>				

Appropriate Care and Associated Codes to Bill			
	CPT and CPT II	ICD-10	Prescription
<p><b>Disease-Modifying Anti-Rheumatic Drug Therapy for Rheumatoid Arthritis (ART) (Admin)</b></p> <p>The percentage of members age 18 years and older who were diagnosed with rheumatoid arthritis and who were dispensed at least one ambulatory prescription for a disease modifying anti-rheumatic drug (DMARD).</p>		<p><b>Rheumatoid Arthritis</b> M05.00, M05.011, M05.012, M05.019, M05.021, M05.022, M05.029, M05.031, M05.032, M05.039, M05.041, M05.042, M05.049, M05.051, M05.052, M05.059, M05.061, M05.062, M05.069, M05.071, M05.072, M05.079, M05.09, M05.10, M05.111, M05.112, M05.119, M05.121, M05.129, M05.131, M05.132, M05.139, M05.141, M05.142, M05.149, M05.151, M05.152, M05.159, M05.161, M05.162, M05.169, M05.171, M05.172, M05.179, M05.19, M05.20, M05.211, M05.212, M05.219, M05.221, M05.222, M05.229, M05.231, M05.232, M05.239, M05.241, M05.242, M05.249, M05.251, M05.252, M05.259, M05.261, M05.262, M05.269, M05.271, M05.272, M05.279, M05.29, M05.30, M05.311, M05.312, M05.319, M05.321, M05.322, M05.329, M05.331, M05.332, M05.339, M05.341, M05.342, M05.349, M05.351, M05.352, M05.359, M05.361, M05.362, M05.369, M05.371, M05.372, M05.379, M05.39, M05.40, M05.411, M05.412, M05.419, M05.421, M05.422, M05.429, M05.431, M05.432, M05.439, M05.441, M05.442, M05.449, M05.451, M05.452, M05.459, M05.461, M05.462, M05.469, M05.471, M05.472, M05.479, M05.49, M05.50, M05.511, M05.512, M05.519, M05.521, M05.522, M05.529, M05.531, M05.532, M05.539, M05.541, M05.542, M05.549, M05.551, M05.552, M05.559, M05.561, M05.562, M05.569, M05.571, M05.572, M05.579, M05.59, M05.60, M05.611, M05.612, M05.619, M05.621, M05.622, M05.629, M05.631, M05.632, M05.639, M05.641, M05.642, M05.649, M05.651, M05.652, M05.659, M05.661, M05.662, M05.669, M05.671, M05.672, M05.679, M05.69, M05.70, M05.711, M05.712, M05.719, M05.721, M05.722, M05.729, M05.731, M05.732, M05.739, M05.741, M05.742, M05.749, M05.751, M05.752, M05.759, M05.761, M05.762, M05.769, M05.771, M05.772, M05.779, M05.79, M05.80, M05.811, M05.812, M05.819, M05.821, M05.822, M05.829, M05.831, M05.832, M05.839, M05.841, M05.842, M05.849, M05.851, M05.852, M05.859, M05.861, M05.862, M05.869, M05.871, M05.872, M05.879, M05.89, M05.9, M06.00, M06.011, M06.012, M06.019, M06.021, M06.022, M06.029, M06.031, M06.032, M06.039, M06.041, M06.042, M06.049, M06.051, M06.052, M06.059, M06.061, M06.062, M06.069, M06.071, M06.072, M06.079, M06.08, M06.09, M06.1, M06.20, M06.211, M06.212, M06.219, M06.221, M06.222, M06.229, M06.231, M06.232, M06.239, M06.241, M06.242, M06.249, M06.251, M06.252, M06.259, M06.261, M06.262, M06.269, M06.271, M06.272, M06.279, M06.28, M06.29</p> <p><i>Continued on page 11</i></p>	

Appropriate Care and Associated Codes to Bill		
CPT and CPT II	ICD-10	Prescription
<p><i>Disease-Modifying Anti-Rheumatic Drug Therapy for Rheumatoid Arthritis (ART) (Admin) continued from page 10</i></p>	<p><b>Rheumatoid Arthritis continued from page 10</b>                      M06.30, M06.311, M06.312, M06.319, M06.321, M06.322, M06.329, M06.331, M06.332, M06.339, M06.341, M06.342, M06.349, M06.351, M06.352, M06.359, M06.361, M06.362, M06.369, M06.371, M06.372, M06.379, M06.38, M06.39, M06.80, M06.811, M06.812, M06.819, M06.821, M06.822, M06.829, M06.831, M06.832, M06.839, M06.841, M06.842, M06.849, M06.851, M06.852, M06.859, M06.861, M06.862, M06.869, M06.871, M06.872, M06.879, M06.88, M06.89, M06.9</p>	
<p><b>Pharmacotherapy Management of COPD Exacerbation (PCE) (Admin)</b></p> <p>The percentage of COPD exacerbations for members 40 years of age and older who had an acute inpatient discharge or ED visit on or between January 1 and November 30 of the measurement year, and who were dispensed appropriate medications.</p> <ol style="list-style-type: none"> <li>1. Dispensed a systemic corticosteroid (or there was evidence of an active prescription) within 14 days of the event.</li> <li>2. Dispensed a bronchodilator (or there was evidence of an active prescription) within 30 days of the event.</li> </ol>	<p><b>Chronic Bronchitis</b>                      J41.0, J41.1, J41.8, J42</p> <p><b>Emphysema</b>                      J43.0, J43.1, J43.2, J43.8, J43.9</p> <p><b>COPD</b>                      J44.0, J44.1, J44.9</p>	<p><b>Glucocorticoids</b>                      Betamethasone, Cortisone-acetate, Dexamethasone, Hydrocortisone, Methylprednisolone, Prednisolone, Prednisone, Triamcinolone</p> <p><b>Bronchodilator Medications</b></p> <p><b>Anticholinergic Agents</b>                      Albuterol-ipratropium, Acclidinium-bromide, Ipratropium, Tiotropium, Umeclidinium</p> <p><b>Beta 2-Agonists</b>                      Albuterol, Arformoterol, Budesonide-formoterol, Fluticasone-salmeterol, Fluticasone-vilanterol, Formoterol, Formoterol-glycopyrrolate, Indacaterol, Indacaterol-glycopyrrolate, Levalbuterol, Mometasone-formoterol, Metaproterenol, Olodaterol hydrochloride, Olodaterol-tiotropium, Pirbuterol, Salmeterol, Umeclidinium-vilanterol</p> <p><b>Methylxanthines</b>                      Aminophylline, Dyphylline, Theophylline</p> <p><b>Antiasthmatic Combinations</b>                      Dyphylline-guaifenesin, Guaifenesin-theophylline</p>

Appropriate Care and Associated Codes to Bill				
	CPT and CPT II	ICD-10	HCPCS	
<p><b>Comprehensive Diabetes Care (CDC) (Hybrid)</b></p> <p>The percentage of members 18–75 years of age with diabetes (type 1 and type 2) who had one of each of the following screenings:</p> <ul style="list-style-type: none"> <li>• Hemoglobin A1C (HbA1C) Testing</li> <li>• Eye Exam (Retinal) Performed</li> <li>• Blood Pressure</li> </ul>	<p><b>Hemoglobin A1C (HbA1C) Testing</b></p> <p><b>HbA1C Screening (CPT)</b> 83036, 83037</p> <p><b>HbA1C Results (CPT II Codes)</b> 3044F (&lt;7.0%) 3045F (7.0–9.0%) 3046F (≥9.0%)</p>	<p><b>Diabetes Diagnoses</b></p> <p>E10.10, E10.11, E10.21, E10.22, E10.29, E10.311, E10.319, E10.321, E10.3211, E10.3212, E10.3213, E10.3219, E10.329, E310.3291, E310.3292, E10.3293, E310.3299, E10.331, E10.3311, E10.3312, E10.3313, E10.3319, E10.339, E10.3391, E10.3392, E10.3393, E10.3399, E10.341, E10.3411, E10.3412, E10.3413, E10.3419, E10.349, E10.3491, E10.3492, E10.3493, E10.3499, E10.351, E10.3511, E10.3512, E10.3513, E10.3519, E10.3521, E10.3522, E103523, E10.3529, E10.3531, E10.3532, E103533, E10.3539, E10.3541, E10.3542, E10.3543, E10.3549, E10.3551, E10.3552, E10.3553, E10.3559, E10.359, E10.3591, E10.3592, E10.3599, E10.36, E10.37X1, E10.37X2, E10.37X3, E10.37X9, E10.39, E10.40, E10.41, E10.42, E10.43, E10.44, E10.49, E10.51, E10.52, E10.59, E10.610, E10.618, E10.620, E10.621, E10.622, E10.628, E10.630, E10.638, E10.641, E10.649, E10.65, E10.69, E10.8, E10.9, E11.00, E11.01, E11.10, E11.11, E11.21, E11.22, E11.29, E11.311, E11.319, E11.321, E11.329, E11.3211, E11.3212, E11.3213, E11.3219, E11.3291, E11.3292, E11.3293, E11.3299, E11.331, E11.3311, E11.3312, E11.3313, E11.3319, E11.339, E11.3391, E11.3392, E11.3393, E11.3399, E11.341, E11.3411, E11.3412, E11.3413, E11.3419, E11.349, E11.351, E11.359, E11.36, E11.39, E11.3411, E11.3412, E11.3413, E11.3419, E11.3491, E11.3492, E11.3493, E11.3499, E11.3511, E11.3512, E11.3513, E11.3519, E11.3521, E11.3522, E11.3523, E11.3529, E11.3531, E113532, E11.3533, E11.3539, E11.3541, E11.3542, E11.3543, E11.3549, E11.3551, E11.3552, E11.3553, E11.3559, E11.3591, E11.3592, E11.3593, E11.3599, E11.37X1, E11.37X2, E11.37X3, E11.37X9, E11.40, E11.41, E11.42, E11.43, E11.44, E11.49, E11.51, E11.52, E11.59, E11.610, E11.618, E11.620, E11.621, E11.622, E11.628, E11.630, E11.638, E11.641, E11.649, E11.65, E11.69, E11.8, E11.9, E13.00, E13.01, E13.10, E13.11, E13.21, E13.22, E13.29, E13.311, E13.319, E13.321, E13.3211, E13.3212, E13.3213, E13.3219, E13.329, E13.3291, E13.3292, E13.3293, E13.3299, E13.331, E13.3311, E13.3312, E13.3313, E13.3319, E13.339, E13.3391, E13.3392, E13.3393, E13.3399, E13.341, E13.3411, E13.3412, E13.3413, E13.3419, E13.349, E13.3491, E13.3492, E13.3493, E13.3499, E13.351, E13.3511, E13.3512, E13.3513, E13.3519, E13.3519, E13.3521, E13.3522, E13.3523, E13.3529, E13.3531, E13.3532, E13.3533, E13.3539, E13.3541, E13.3542, E13.3543, E13.3549, E13.3551, E13.3552, E13.3553, E13.3559, E13.359, E13.3591, E13.3592, E13.3593, E13.3599, E13.36, E13.37X1, E13.37X2, E1337X3, E13.37X9, E13.39, E13.40, E13.41, E13.42, E13.43, E13.44, E13.49, E13.51, E13.52, E13.59, E13.610, E13.618, E13.620, E13.621, E13.622, E13.628, E13.630, E13.638, E13.641, E13.649, E13.65, E13.69, E13.8, E13.9</p> <p>O24.011, O24.012, O24.013, O24.019, O24.02, O24.03, O24.111, O24.112, O24.113, O24.119, O24.12, O24.13, O24.311, O24.312, O24.313, O24.319, O24.32, O24.33, O24.811, O24.812, O24.813, O24.819, O24.82, O24.83</p>	<p><b>Eye Exam (Retinal) Performed</b></p> <p>S0620, S0621, S3000</p>	
	<p><b>Eye Exam (Retinal) Performed</b></p> <p><b>CPT Codes</b></p> <p><b>Diabetic Retinal Screening</b> 67028, 67030, 67031, 67036, 67039, 67040, 67041, 67042, 67043, 67101, 67105, 67107, 67108, 67110, 67113, 67121, 67141, 67145, 67208, 67210, 67218, 67220, 67221, 67227, 67228, 92002, 92004, 92012, 92014, 92018, 92019, 92134, 92225, 92226, 92227, 92228, 92230, 92235, 92240, 92250, 92260, 99203, 99204, 99205, 99213, 99214, 99215, 99242, 99243, 99244, 99245</p> <p><b>Eye Enucleation</b> 65091, 65093, 65101, 65103, 65105, 65110, 65112, 65114</p> <p><b>CPT II Codes</b></p> <p><b>Dilated Retinal Eye Exam with Interpretation</b> 2022F</p> <p><b>7 Standard Field Stereoscopic Photos with Interpretation</b> 2024F</p> <p><b>Eye Imaging Validated</b> 2026F</p> <p><b>Low Risk for Retinopathy (no evidence of retinopathy in the prior year)</b> 3072F</p>			<p><b>Blood Pressure</b></p> <p><b>Systolic CPT Codes</b> 3074F (&lt;140 mm Hg) 3075F (130–139 mm/Hg) 3077F (≥140 mm/Hg)</p> <p><b>Diastolic CPT Codes</b> 3078F (&lt;80 mm Hg) 3079F (80–89 mm Hg) 3080F (≥90 mm Hg)</p> <p><i>Please record the exact reading in the medical record and do not round the sphygmomanometer result.</i></p>

	Appropriate Care and Associated Codes to Bill			
	CPT and CPT II	CPT-CAT-11	ICD-10-CM Diagnosis	UB Revenue
Medical Attention for Nephropathy Urine Protein Test	Urine Macroalbin Tests 81000, 81001, 81002, 81003, 81005, 82042, 84043, 84044, 84156	Positive Microalbuminuria Test Result Documented and Reviewed (DM) 3060F, 3062F  Negative Microalbuminuria Test Result Documented and Reviewed (DM) 3061F		
Medical Attention for Nephropathy Nephropathy Treatment	Evidence of ESRD 36147, 36800, 36810, 36815, 36818, 36819, 36820, 36821, 36831, 36832, 36833, 90935, 90937, 90940, 90945, 90947, 90951, 90952, 90953, 90954, 90955, 90956, 90957, 90958, 90959, 90960, 90961, 90962, 90963, 90964, 90965, 90966, 90967, 90968, 90969, 90970, 90989, 90993, 90997, 90999, 99512  Nephropathy Screening Tests 81000, 81001, 81002, 81003, 81005, 82042, 82043, 82044, 84156	Nephropathy Screening Test 3060F, 3061F  Treatment for Nephropathy 3066F, 4010F	E08.21, E08.22, E08.29, E09.21, E09.22, E09.29, E10.21, E10.22, E10.29, E11.21, E11.22, E11.29, E13.21, E13.22, E13.29 I12.0, I12.9, I13.0, I13.10, I13.11, I13.2, I15.0, I15.1 N00.0, N00.1, N00.2, N00.3, N00.4, N00.5, N00.6, N00.7, N00.8, N00.9, N01.0, N01.1, N01.2, N01.3, N01.4, N01.5, N01.6, N01.7, N01.8, N01.9, N02.0, N02.1, N02.2, N02.3, N02.4, N02.5, N02.6, N02.7, N02.8, N02.9, N03.0, N03.1, N03.2, N03.3, N03.4, N03.5, N03.6, N03.7, N03.8, N03.9, N04.0, N04.1, N04.2, N04.3, N04.4, N04.5, N04.6, N04.7, N04.8, N04.9, N05.0, N05.1, N05.2, N05.3, N05.4, N05.5, N05.6, N05.7, N05.8, N05.9, N06.0, N06.1, N06.2, N06.3, N06.4, N06.5, N06.6, N06.7, N06.8, N06.9, N07.0, N07.1, N07.2, N07.3, N07.4, N07.5, N07.6, N07.7, N07.8, N07.9, N08, N14.0, N14.1, N14.2, N14.3, N14.4, N17.0, N17.1, N17.2, N17.8, N17.9, N18.1, N18.2, N18.3, N18.4, N18.5, N18.6, N18.9, N19, N25.0, N25.1, N25.81, N25.89, N25.9, N26.1, N26.2, N26.9 Q60.0, Q60.1, Q60.2, Q60.3, Q60.4, Q60.5, Q60.6, Q61.00, Q61.01, Q61.02, Q61.11, Q61.19, Q61.2, Q61.3, Q61.4, Q61.5, Q61.8, Q61.9 R80.0, R80.1, R80.2, R80.3, R80.8, R80.9	0800, 0801, 0802, 0803, 0804, 0809, 0820, 0821, 0822, 0823, 8024, 0825, 0829, 0830, 0831, 0832, 0833, 0834, 0835, 0839, 0840, 0841, 0842, 0843, 0844, 0845, 0849, 0850, 0851, 0852, 0853, 0854, 0855, 0859, 0880, 0881, 0882, 0889



Appropriate Care and Associated Codes to Bill		
CPT and CPT II	HCPCS	Prescription
<p><b>Annual Monitoring for Patients on Persistent Medications (MPM) (Admin)</b></p> <p>The percentage of members 18 years of age and older who received at least 180 treatment days of ambulatory medication therapy for a select therapeutic agent during the measurement year and at least one therapeutic monitoring event for the therapeutic agent in the measurement year.</p> <ul style="list-style-type: none"> <li>Annual monitoring for members on angiotensin converting enzymes (ACE) inhibitors or angiotensin receptor blockers (ARB)</li> <li>Annual monitoring for members on diuretics</li> <li>Total rate (the sum of the two numerators divided by the sum of the two denominators)</li> </ul>	<p><b>Physiologic Monitoring Tests</b> 80047, 80048, 80050, 80053, 80069 (Lab panel)</p> <p><b>Serum Potassium</b> 80051, 84132</p> <p><b>Serum Creatinine</b> 82565, 82575</p>	<p><b>ACE/ARB Medications</b></p> <p><b>Angiotensin Converting Enzyme Inhibitors</b> Benazepril, Captopril, Enalapril, Fosinopril, Lisinopril, Moexipril, Perindopril, Quinapril, Ramipril, Trandolapril</p> <p><b>Angiotensin II Inhibitors</b> Azilsartan, Candesartan, Eprosartan, Irbesartan, Losartan, Olmesartan, Telmisartan, Valsartan</p> <p><b>Antihypertensive Combinations</b> Aliskiren-valsartan, Amlodipine-benazepril, Amlodipine-hydrochlorothiazide-valsartan, Amlodipine-hydrochlorothiazide-olmesartan, Amlodipine-olmesartan, Amlodipine-perindopril, Amlodipine-telmisartan, Amlodipine-valsartan, Azilsartan-chlorthalidone, Benazepril-hydrochlorothiazide, Candesartan-hydrochlorothiazide, Captopril-hydrochlorothiazide, Enalapril-hydrochlorothiazide, Eprosartan-hydrochlorothiazide, Fosinopril-hydrochlorothiazide, Hydrochlorothiazide-irbesartan, Hydrochlorothiazide-lisinopril, Hydrochlorothiazide-losartan, Hydrochlorothiazide-moexipril, Hydrochlorothiazide-olmesartan, Hydrochlorothiazide-quinapril, Hydrochlorothiazide-telmisartan, Hydrochlorothiazide-valsartan, Sacubitril-valsartan, Trandolapril-verapamil</p> <p><b>Diuretic Medications</b></p> <p><b>Antihypertensive Combinations</b> Aliskiren-hydrochlorothiazide, Aliskiren-hydrochlorothiazide-amlodipine, Amiloride-hydrochlorothiazide, Amlodipine-hydrochlorothiazide-olmesartan, Amlodipine-hydrochlorothiazide-valsartan, Atenolol-chlorthalidone, Azilsartan-chlorthalidone, Benazepril-hydrochlorothiazide, Bendroflumethiazide-nadolol, Bisoprolol-hydrochlorothiazide, Candesartan-hydrochlorothiazide, Captopril-hydrochlorothiazide, Chlorthalidone-clonidine, Enalapril-hydrochlorothiazide, Eprosartan-hydrochlorothiazide, Fosinopril-hydrochlorothiazide, Hydrochlorothiazide-irbesartan, Hydrochlorothiazide-lisinopril, Hydrochlorothiazide-losartan, Hydrochlorothiazide-methyldopa, Hydrochlorothiazide-metoprolol, Hydrochlorothiazide-moexipril, Hydrochlorothiazide-olmesartan, Hydrochlorothiazide-propranolol, Hydrochlorothiazide-quinapril, Hydrochlorothiazide-spirolactone, Hydrochlorothiazide-telmisartan, Hydrochlorothiazide-triamterene, Hydrochlorothiazide-valsartan</p> <p><b>Loop Diuretics</b> Bumetanide, Ethacrynic acid, Furosemide, Torsemide</p> <p><b>Potassium-Sparing Diuretics</b> Amiloride, Eplerenone, Spirolactone, Triamterene Thiazide diuretics: Chlorothiazide, Chlorthalidone, Hydrochlorothiazide, Indapamide, Methyclothiazide, Metolazone</p>

	Appropriate Care and Associated Codes to Bill		
	CPT and CPT II	HCPCS	Prescription
<p><b>Persistence of Beta-Blocker Treatment After a Heart Attack (PBH) (Admin)</b></p> <p>The percentage of members 18 years and older during the measurement year who were hospitalized and discharged from July 1 of the year prior to the measurement year to June 30 of the measurement year with a diagnosis of AMI and who received persistent beta-blocker treatment for six months after discharge.</p>		<p><b>Acute MI</b> I21.01, I21.02, I21.09, I21.11, I21.19, I21.21, I21.29, I21.3, I21.4</p> <p><b>Acute Myocardial Infarction Unspecified</b> I21.9</p> <p><b>Myocardial Infarction Type 2</b> I21.A1</p> <p><b>Other Myocardial Infarction Type</b> I21.A9</p>	<p><b>PBH Beta-Blocker Medications</b> Carvedilol, Labetalol, Nadolol, Penbutolol, Pindolol, Propranolol, Timolol, Sotalol, Acebutolol, Atenolol, Betaxolol, Bisoprolol, Metoprolol, Nebivolol, Atenolol-chlorthalidone, Bendroflumethiazide-nadolol, Bisoprolol-hydrochlorothiazide, Hydrochlorothiazide-metoprolol, Hydrochlorothiazide-propranolol</p>