



2018 Behavioral Health Coding Reference Guide for Providers



Contents

| | |
|--|-------|
| Follow-up Care for Children Prescribed ADHD Medication (ADD) | 3 |
| Use of Multiple Concurrent Antipsychotics in Children and Adolescents (APC)..... | 4 |
| Metabolic Monitoring for Children and Adolescents on Antipsychotics (APM) | 4 |
| Use of First-Line Psychosocial Care for Children and Adolescents on Antipsychotics (APP) | 5 |
| Anti-depressant Medication Management (AMM) | 5 |
| Follow-up after Emergency Department Visit for Alcohol and other Drug Dependence (FUA) | 6 |
| Follow-up after Emergency Department Visit for Alcohol and other Drug Dependence (FUA) | 7 |
| Follow-up after Hospitalization for Mental Illness (FUH) | 8 |
| Follow-up after Hospitalization for Mental Illness (FUH) | 9 |
| Follow-up after Emergency Department visits for Mental Illness (FUM) | 10 |
| Follow-up after Emergency Department visits for Mental Illness (FUM) | 11 |
| Initiation and Engagement of Alcohol and Other Drug Dependence (AOD) Treatment (IET)..... | 12 |
| Initiation and Engagement of Alcohol and Other Drug Dependence (AOD) Treatment (IET)..... | 13 |
| Adherence to Antipsychotic Medications for Individuals with Schizophrenia (SAA)..... | 14 |
| Adherence to Antipsychotic Medications for Individuals with Schizophrenia (SAA)..... | 15 |
| Cardiovascular Monitoring for People with Cardiovascular Disease and Schizophrenia (SMC) | 16–19 |
| Diabetes Monitoring for People with Diabetes and Schizophrenia (SMD) | 20–22 |
| Diabetes Screening for People with Schizophrenia or Bipolar Disorder who are using Antipsychotic Medications (SSD) | 23–25 |

Current Procedural Terminology ©2014 American Medical Association. All rights reserved.

Current Procedural Terminology (CPT) and Uniform Bill (UB) codes are stated to the minimum specificity required.

2018 Behavioral Health Guide

| Appropriate Care and Associated Codes to Bill | | | | |
|--|---|--------------|--|--|
| | CPT Codes | ICD-10 Codes | HCPCS Codes | UB Revenue |
| <p>Follow-up Care for Children Prescribed ADHD Medication (ADD)</p> <p>The percentage of children (6–12 years of age) who are newly prescribed attention-deficit/hyperactivity disorder (ADHD) medication and had at least 3 follow-up visits. Two phases are reported:</p> <p>The first is the Initiation Phase in which the child is seen by a prescribing practitioner within 30 days after the prescription start date but not on the day the script is filled.</p> <p>The second phase is the continuation & maintenance phase. This is where the child needs 2 follow-up visits on different dates of service with any practitioner between day 31–300 after prescription start date.</p> | <p>ADD Stand Alone Visits Value Set, use with UBREV and HCPCS codes as appropriate for both Initiation and Continuation & Maintenance Phase</p> <p>96150–96154, 98960–98962, 99078, 99201–99205, 99211–99215, 99217–99220, 99241–99245, 99341–99345, 99347–99350, 99383, 99384, 99393, 99394, 99401–99404, 99411, 99412, 99510</p> <p>Use for both Initiation and Continuation & Maintenance Phase the ADD Visits Group 1 Value Sets with ADD POS Group 1 Value Set:</p> <p>90791, 90792, 90832–90834, 90836–90840, 90845, 90847, 90849, 90853, 90875, 90876</p> <p>POS Group 1 Value Set use with ADD Visits Value Set for both phases:</p> <p>03, 05, 07, 09–20, 22, 33, 49, 50, 52, 53, 71, 72</p> <p>ADD Visits Group 2 Value Set Codes used with the ADD POS Group 2 Value Set for both phases:</p> <p>99221–99223, 99231–99233, 99238, 99239, 99251–99255</p> <p>ADD POS Group 2 Value Set use with ADD Visits Group 2 Value Set for both phases:</p> <p>52, 53</p> <p>Do not count visits billed with a telehealth Modifier Value Set or a telehealth POS Value Set in the Initiation Phase.</p> <p>In the Continuation & Maintenance Phase can use one of the two visits as a telephone visit with the following codes:</p> <p>98966–98968, 99441–99443</p> <p>For the Continuation & Maintenance Phase can use the ADD Visits Group 1 or 2 Value Set with Telehealth POS Value Set. (Only 1 visit out of two can be either telephone or telehealth): Use code: GT</p> | | <p>Use with the Initiation and the Continuation & Maintenance phase codes with the ADD Stand Alone Visits Value Set Codes:</p> <p>G0176, G0177, G0410, G0411, G0463, H0004, H0031, H0034–H0037, H0039, H0040, H2000, H2001, H2010–H2020, M0064, S0201, S9480, S9484, S9485, T1015</p> | <p>In the Initiation & Continuation & Maintenance Phase use the following UB Rev codes with the ADD Stand Alone visits Value Set:</p> <p>0510, 0513, 0515, 0516, 0517, 0519–0523, 0526–0529, 0900, 0902–0905, 0907, 0911–0917, 0919, 0982, 0983</p> |

2018 Behavioral Health Guide

| | Appropriate Care and Associated Codes to Bill | | | |
|---|---|--------------|-------------|------------|
| | CPT Codes | ICD-10 Codes | HCPCS Codes | UB Revenue |
| <p>Use of Multiple Concurrent Antipsychotics in Children and Adolescents (APC)</p> <p>Children and adolescents 1–17 years of age who were on two or more concurrent antipsychotic medications.</p> | | | | |
| <p>Metabolic Monitoring for Children and Adolescents on Antipsychotics (APM)</p> <p>Children and adolescents 1–17 years of age who had two or more antipsychotic prescriptions and had metabolic testing.</p> <p>Note: To qualify you need at least two antipsychotic medication dispensing events of the same or different medications on different dates of service during the measurement year.</p> <p>Note: Member needs to have both a blood glucose or HbA1c and a LDL-C or Cholesterol blood testing on the same or different dates of service during the measurement year.</p> | <p>Metabolic Testing:</p> <p>Cholesterol Tests Other Than LDL 82465, 83718, 84478, 2085–9, 2093–3, 2571–8, 5932–9</p> <p>Glucose Tests 80047, 80048, 80050, 80053, 80069, 82947, 82950, 82951, 10450–5, 14753–8, 14754–6, 14756–1, 14757–9, 14759–5, 14764–5, 14765–2, 1492–8, 1494–4, 1496–9, 1499–3, 14995–5, 1501–6, 1504–0, 1507–3, 1514–9, 1518–0, 1530–5, 1533–9, 1554–5, 1557–8, 1558–6, 17865–7, 20436–2, 20437–0, 20438–8, 25666–9, 26554–6, 30251–3, 30265–3, 30267–9, 32320–4, 40285–9, 40286–7, 41024–1, 49134–0, 51597–3, 55351–1, 55381–8, 6749–6, 74774–1, 9375–7</p> <p>HbA1c 83036, 83037, 3044F–3046F, 17856–6, 4548–4, 4549–2</p> <p>LDL-C Tests 80061, 83700, 83701, 83704, 83721, 3048F–3050F, 12773–8, 13457–7, 18261–8, 18262–6, 2089–1, 49132–4, 55440–2</p> | | | |

2018 Behavioral Health Guide

| | Appropriate Care and Associated Codes to Bill | | | |
|--|--|---|---|---|
| | CPT Codes | ICD-10 Codes | HCPCS Codes | UB Revenue |
| <p>Use of First-Line Psychosocial Care for Children and Adolescents on Antipsychotics (APP)</p> <p>Members of children and adolescents 1–17 years of age who had a new prescription for an antipsychotic medication and had documentation of psychosocial care as first-line treatment.</p> | <p>CPT Codes for Psychosocial Care Value Set with or without telehealth modifier:</p> <p>90832–90834, 90836–90840, 90845–90847, 90849, 90853, 90875, 90876, 90880</p> <p>Telehealth Modifier Code:</p> <p>95, GT</p> | | <p>HCPCS Codes for Psychosocial Care Value Set with or without Telehealth Modifier:</p> <p>G0176, G0177, G0409–G0411, H0004, H0035–H0040, H2000, H2001, H2011–H2014, H2017–H2020, S0201, S9480, S9484, S9485</p> | |
| <p>Anti-depressant Medication Management (AMM)</p> <p>Members 18 years of age and older who were treated with antidepressant medication, had a diagnosis of major depression and who remained on an antidepressant medication treatment for 84 to 180 days (3 to 6 months).</p> | <p>AMM Stand Alone Visits Value Set with Major Depression Value Set with or without telehealth modifier value set.</p> <p>98960–98962, 99078, 99201–99205, 99211–99215, 99217–99220, 99241–99245, 99341–99345, 99347–99350, 99384–99387, 99394–99397, 99401–99404, 99411, 99412, 99510</p> <p>AMM visit with Place of Service (POS) Codes and with Major Depression with or without Telehealth Modifier Value.</p> <p>90791, 90792, 90832–90834, 90836–90840, 90845, 90847, 90849, 90853, 90867–90870, 90875, 90876, 99221–99223, 99231–99233, 99238, 99239, 99251–99255</p> <p>POS Codes:</p> <p>3, 5, 7, 9, 11–20, 22, 24, 33, 49, 50, 52, 53, 71, 72</p> <p>Telehealth Modifier Value Set:</p> <p>95, GT</p> <p>Telephone Visits Value Set with Major Depression Value Set:</p> <p>98966–98968, 99441–99443.</p> | <p>Major Depression Disorder Value Set:</p> <p>F32.0–F32.9, F33.0–F33.3, F33.41, F33.9</p> | <p>AMM Stand Alone visits HCPCS codes.</p> <p>G0155, G0176, G0177, G0409–G0411, G0463, H0002, H0004, H0031, H0034–H0037, H0039, H0040, H2000, H2001, H2010–H2020, M0064, S0201, S9480, S9484, T1015</p> | <p>AMM Stand Alone Visits UB Revenue Codes:</p> <p>0510, 0513, 0515–0517, 0519–0523, 0526–0529, 0900–0905, 0907, 0911–0917, 0919, 0982, 0983</p> |

2018 Behavioral Health Guide

| | Appropriate Care and Associated Codes to Bill | | | |
|--|--|--|--|------------|
| | CPT Codes | ICD-10 Codes | HCPCS Codes | UB Revenue |
| <p>Follow-up after Emergency Department Visit for Alcohol and other Drug Dependence (FUA)</p> <p>The percentage of emergency department (ED) visits for members 13 years of age and older with a principal diagnosis of alcohol or other drug (AOD) abuse or dependence, who had a follow up visit for AOD. Two rates are reported: 1) Percentage of ED visits for which the member received follow-up within 30 days of the ED visit. 2) Received follow-up within 7 days of the ED visit.</p> | <p>IET stand alone visits value set with a principal diagnosis of AOD abuse or dependence with or without telehealth modifier value sets.</p> <p>98960–98962, 99078, 99201–99205, 99211–99215, 99217–99220, 99241–99245, 99341–99345, 99347–99350, 99384–99387, 99394–99397, 99401–99404, 99408, 99409, 99411, 99412, 99510</p> <p>IET Visits Group 1 Value Set with IET POS Group 1 Value Set and a principal diagnosis of AOD abuse or dependence with or without a telehealth modifier.</p> <p>90791, 90792, 90832–90834, 908836–90840, 908845, 90847, 90849, 90853, 90875, 90876</p> <p>IET POS Group 1 Value Set</p> <p>3, 5, 7, 9, 11–20, 22, 33, 49, 50, 52, 53, 57, 71, 72</p> <p>Telehealth Modifier Value Set:</p> <p>95, GT</p> <p>IET visits Group 2 Value Set with IET POS Group 2 Value Set and a principal diagnosis of AOD abuse or dependence with or without a telehealth modifier</p> <p><i>(Additional CPT codes on page 7)</i></p> | <p>Alcohol and Other Drug Dependence Codes:</p> <p>F10.10, F10.120, F10.121, F10.129, F10.14, F10.150, F10.151, F10.159, F10.180, F10.181, F10.182, F10.188, F10.19, F10.20, F10.220, F10.221, F10.229, F10.230, F10.231, F10.232, F10.239, F10.24, F10.250, F10.251, F10.259, F10.26, F10.27, F10.280, F10.281, F10.282, F10.288, F10.29, F11.10, F11.120, F11.121, F11.122, F11.129, F11.14, F11.150, F11.151, F11.159, F11.181, F11.182, F11.188, F11.19, F11.20, F11.220, F11.221, F11.222, F11.229, F11.23, F11.24, F11.250, F11.251, F11.259, F11.281, F11.282, F11.288, F11.29, F12.10, F12.120, F12.121, F12.122, F12.129, F12.150, F12.151, F12.159, F12.180, F12.188, F12.19, F12.20, F12.220, F12.221, F12.222, F12.229, F12.250, F12.251, F12.259, F12.280, F12.288, F12.29, F13.10, F13.120, F13.121, F13.129, F13.14, F13.150, F13.151, F13.159, F13.180, F13.181, F13.182, F13.188, F13.19, F13.20, F13.220, F13.221, F13.229, F13.230, F13.231, F13.232, F13.239, F13.24, F13.250, F13.251, F13.259, F13.26, F13.27, F13.280, F13.281, F13.282, F13.288, F13.29, F14.10, F14.120, F14.121, F14.122, F14.129, F14.150, F14.151, F14.159, F14.180, F14.181, F14.182, F14.188, F14.19, F14.20, F14.220, F14.221, F14.222, F14.229, F14.23, F14.24, F14.250, F14.251, F14.259, F14.280, F14.281, F14.282, F14.288, F14.29, F15.10, F15.120, F15.121, F15.122, F15.129, F15.14, F15.150, F15.151, F15.159, F15.180, F15.181, F15.182, F15.188, F15.19, F15.20, F15.220, F15.221, F15.222, F15.229, F15.23, F15.24, F15.250, F15.251, F15.259, F15.280, F15.281, F15.282, F15.288, F15.29, F16.10, F16.120, F16.121, F16.122, F16.129, F16.14, F16.150, F16.151, F16.159, F16.180, F16.183, F16.188</p> <p><i>(Additional ICD-10 codes on page 7)</i></p> | <p>HCPCS to be used with Stand Alone Visit Codes:</p> <p>G0155, G0176, G0177, G0396, G0397, G0409–G0411, G0443, G0463, H0001, H0002, H0004, H0005, H0007, H0015, H0016, H0022, H0031, H0034–H0037, H0039, H0040, H0047, H2000, H2001, H2010–H2020, H2035, H2036, M0064, S0201, S9480, S9484, S9458, T1006, T1012, T1015</p> | |

2018 Behavioral Health Guide

| | Appropriate Care and Associated Codes to Bill | | | |
|--|--|--|-------------|------------|
| | CPT Codes | ICD-10 Codes | HCPCS Codes | UB Revenue |
| Follow-up after Emergency Department Visit for Alcohol and other Drug Dependence (FUA) (Continued from page 6) | IET Visits Group 2 codes: 99221–99223, 99231–99233, 99238, 99239, 99251–99255 IET POS Group 2 Value Set: 52, 53 Telephone Visits Value Set with principal diagnosis of AOD abuse or dependence with or without a telehealth modifier. 98966–98968, 99441–99443 Online Assessments Value Set with a principal diagnosis of AOD abuse or dependence. 98969, 99444 | F16.19, F16.20, F16.220, F16.221, F16.229, F16.24, F16.250, F16.251, F16.259, F16.280, F16.283, F16.288, F16.29, F18.10, F18.120, F18.121, F18.129, F18.14, F18.150, F18.151, F18.159, F18.17, F18.180, F18.188, F18.19, F18.20, F18.220, F18.221, F18.229, F18.24, F18.250, F18.251, F18.259, F18.27, F18.280, F18.288, F18.29, F19.10, F19.120, F19.121, F19.122, F19.129, F19.14, F19.150, F19.151, F19.159, F19.16, F19.17, F19.180, F19.181, F19.182, F19.188, F19.19, F19.20, F19.220, F19.221, F19.222, F19.229, F19.230, F19.231, F19.232, F19.239, F19.24, F19.250, F19.251, F19.259, F19.26, F19.27, F19.280, F19.281, F19.282, F19.288, F19.29, | | |

2018 Behavioral Health Guide

| | Appropriate Care and Associated Codes to Bill | | | |
|--|---|---|--|---|
| | CPT Codes | ICD-10 Codes | HCPCS Codes | UB Revenue |
| <p>Follow-up after Hospitalization for Mental Illness (FUH)</p> <p>Members 6 years of age and older who were hospitalized for treatment of selected mental illness diagnoses and who had a follow-up visit with a mental health practitioner within 7 and/or 30 days of hospital discharge.</p> <p>Note: Cannot count the day of hospital discharge.</p> | <p>FUH Stand Alone Visits Value Set with or without Telehealth Modifier Value Set</p> <p>98960, 98961, 98962, 99078, 99201, 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99217, 99218, 99219, 99220, 99241, 99242, 99243, 99244, 99245, 99341, 99342, 99343, 99344, 99345, 99347, 99348, 99349, 99350, 99383, 99384, 99385, 99386, 99387, 99393, 99394, 99395, 99396, 99397, 99401, 99402, 99403, 99404, 99411, 99412, 99510</p> <p>FUH Visits Group 1 Value Set with FUH POS Group 1 Value Set with or without Telehealth Modifier Value Set</p> <p>90791, 90792, 90832, 90833, 90834, 90836, 90837, 90838, 90839, 90840, 90845, 90847, 90849, 90853, 90867, 90868, 90869, 90870, 90875, 90876</p> <p>FUH POS Group 1 Value Set:</p> <p>3, 5, 7, 9, 11–20, 22, 24, 33, 49, 50, 52, 53, 71, 72</p> <p>FUH Visits Group 2 Value Set with FUH POS Group 2 Value Set with or without a Telehealth Modifier Value Set</p> <p>FUH Visits Group 2:</p> <p>99221–99223, 99231–99233, 99238, 99239, 99251–99255</p> <p>FUH POS Group 2 Value Sets:</p> <p>52, 53</p> <p><i>(Additional CPT codes on page 9)</i></p> | <p>F20.0, F20.1, F20.2, F20.3, F20.5, F20.81, F20.89, F20.9, F21, F22, F23, F24, F25.0, F25.1, F25.8, F25.9, F28, F29, F30.10, F30.11, F30.12, F30.13, F30.2, F30.3, F30.4, F30.8, F30.9, F31.0, F31.10, F31.11, F31.12, F31.13, F31.2, F31.30, F31.31, F31.32, F31.4, F31.5, F31.60, F31.61, F31.62, F31.63, F31.64, F31.70, F31.71, F31.72, F31.73, F31.74, F31.75, F31.76, F31.77, F31.78, F31.81, F31.89, F31.9, F32.0, F32.1, F32.2, F32.3, F32.4, F32.5, F32.8, F32.81, F32.89, F32.9, F33.0, F33.1, F33.2, F33.3, F33.40, F33.41, F33.42, F33.8, F33.9, F34.0, F34.1, F34.8, F34.81, F34.89, F34.9, F39, F42, F42.2, F42.3, F42.4, F42.8, F42.9, F43.0, F43.10, F43.11, F43.12, F43.20, F43.21, F43.22, F43.23, F43.24, F43.25, F43.29, F43.8, F43.9, F44.89, F53, F60.0, F60.1, F60.2, F60.3, F60.4, F60.5, F60.6, F60.7, F60.81, F60.89, F60.9, F63.0, F63.1, F63.2, F63.3, F63.81, F63.89</p> <p><i>(Additional ICD-10 codes on page 9)</i></p> | <p>Use for Stand Alone Codes:</p> <p>G0155, G0176, G0177, G0409–G0411, G0463, H0002, H0004, H0031, H0034–H0037, H0039, H0040, H2000, H2001, H2010–H2020, M0064, S0201, S9480, S9484, S9485, T1015</p> | <p>FUH Rev Codes Group 1 in Behavioral health Setting:</p> <p>0513, 0900–0905, 0907, 0911–0917, 0919</p> <p>FUH Rev Codes Group 2 in Non-Behavioral health Setting with a mental health practitioner:</p> <p>0510, 0515–0517, 0519–0523, 0526–0529, 0982, 0983</p> <p>Visit in a non-behavioral healthcare setting (FUH Rev Code Group 2) use Mental Illness Value Set</p> |

2018 Behavioral Health Guide

| | Appropriate Care and Associated Codes to Bill | | | |
|--|--|---|-------------|------------|
| | CPT Codes | ICD-10 Codes | HCPCS Codes | UB Revenue |
| Follow-up after Hospitalization for Mental Illness (FUH) (Continued from page 8) | Telehealth Modifier Value Set: 95, GT Transitional Care Management Services (TCM) 14 day 99495 and Transitional Care Management Services (TCM) 7 day is 99496 | F63.9, F68.10, F68.11, F68.12, F68.13, F68.8, F84.0, F84.2, F84.3, F84.5, F84.8, F84.9, F90.0, F90.1, F90.2, F90.8, F90.9, F91.0, F91.1, F91.2, F91.3, F91.8, F91.9, F93.0, F93.8, F93.9, F94.0, F94.1, F94.2, F94.8, F94.9 | | |

2018 Behavioral Health Guide

| | Appropriate Care and Associated Codes to Bill | | | |
|--|---|--|---|--|
| | CPT Codes | ICD-10 Codes | HCPCS Codes | UB Revenue |
| <p>Follow-up after Emergency Department visits for Mental Illness (FUM)</p> <p>Emergency department visits for members 6 years of age and older with a principal diagnosis of mental illness, who had a follow-up visit for their mental illness, within 7 and/or 30 days.</p> <p>Note: Member must be 6 years or older on the date of the ED visit.</p> | <p>The following criteria is for 7-Day and 30-Day follow-up:</p> <p>ED Codes: 99281–99285</p> <p>FUH Stand Alone Visit Codes: May use stand alone visit codes with a mental health diagnosis value set (ICD-10) with or without a telehealth modifier value set.</p> <p>98960–98962, 99078, 99201–99205, 99211–99215, 99217–99220, 99241–99245, 99341–99345, 99347–99350, 99383–99387, 99393–99397, 99401–99404, 99411, 99412, 99510, G0155, G0176, G0177, G0409–G0411, G463, H0002, H0004, H0031, H0034–H0037, H0039, H0040, H2000, H2001, H2010–H2020, M0064, S0201, S9480, S9485, T1015</p> <p>FUH Visits Group 1 Value Set with FUH POS Group 1 Value Set with a principal diagnosis of mental health disorder with or without telehealth modifier.</p> <p>90791, 90792, 90832–90840, 90845, 90847, 90849, 90853, 90867, 90868, 90869, 90870, 90875, 90876</p> <p>Use the following POS Codes with FUH Visit Group 1 Codes: 03, 05, 07, 09, 11–20, 22, 24, 33, 49, 50, 52, 53, 71, 72</p> <p>Telehealth Modifier Value Set 95, GT</p> <p>Use FUH Visits Group 2 Value Set with FUH POS Group 2 Value Sets with a principal diagnosis of a mental health disorder with or without telehealth modifier.</p> <p><i>(Additional CPT codes on page 11)</i></p> | <p>Mental Health Diagnosis Codes:</p> <p>F03.90, F03.91, F20.0, F20.1, F20.2, F20.3, F20.5, F20.81, F20.89, F20.9, F21, F22, F23, F24, F25.0, F25.1, F25.8, F25.9, F28, F29, F30.10, F30.11, F30.12, F30.13, F30.2, F30.3, F30.4, F30.8, F30.9, F31.0, F31.10, F31.11, F31.12, F31.13, F31.2, F31.30, F31.31, F31.32, F31.4, F31.5, F31.60, F31.61, F31.62, F31.63, F31.64, F31.70, F31.71, F31.72, F31.73, F31.74, F31.75, F31.76, F31.77, F31.78, F31.81, F31.89, F31.9, F32.0, F32.1, F32.2, F32.3, F32.4, F32.5, F32.8, F32.81, F32.89, F32.9, F33.0, F33.1, F33.2, F33.3, F33.40, F33.41, F33.42, F33.8, F33.9, F34.0, F34.1, F34.8, F34.81, F34.89, F34.9, F39, F40.00, F40.01, F40.02, F40.10, F40.11, F40.210, F40.218, F40.220, F40.228, F40.230, F40.231, F40.232, F40.233, F40.240, F40.241, F40.242, F40.243, F40.248, F40.290, F40.291, F40.298, F40.8, F40.9, F41.0, F41.1, F41.3, F41.8, F41.9, F42, F42.2–F42.4, F42.8, F42.9, F43.0, F43.10, F43.11, F43.12, F43.20, F43.21, F43.22, F43.23, F43.24, F43.25, F43.29, F43.8, F43.9, F44.0, F44.1, F44.2, F44.4, F44.5, F44.6, F44.7, F44.81, F44.89, F44.9, F45.0, F45.1, F45.20, F45.21, F45.22, F45.29, F45.41, F45.42, F45.8, F45.9, F48.1, F48.2, F48.8, F48.9, F50.00, F50.01, F50.02, F50.2, F50.8, F50.81, F50.89, F50.9, F51.01, F51.02, F51.03, F51.04, F51.05, F51.09, F51.11, F51.12, F51.13, F51.19, F51.3, F51.4, F51.5, F51.8, F51.9, F52.0, F52.1, F52.21, F52.22, F52.31, F52.32, F52.4, F52.5, F52.6, F52.8, F52.9,</p> <p><i>(Additional ICD-10 codes on page 11)</i></p> | <p>Used with FUH Stand Alone Visits:</p> <p>G0155, G0176, G0177, G0409–G0411, G0463, H0002, H0004, H0031, H0034–H0037, H0039, H0040, H2000, H2001, H2010–H2020, M0064, S0201, S9480, S9484, S9485, T1015</p> | <p>Use FUH REV Codes Group 1 for a visit to a behavioral healthcare setting</p> <p><i>Reminder, no Mental Health Diagnosis Codes with FUH Rev Codes Group 1.</i></p> <p>0513, 0900–0905, 0911–0917, 0919</p> <p>Follow up after ED Visit Codes:</p> <p>0450, 0451, 0452, 0456, 0459</p> <p>For a visit to a nonbehavioral healthcare setting with a principal diagnosis of a mental health disorder use FUH RevCodes Group 2 with the mental health diagnosis value set:</p> <p>0510, 0515–0517, 0519–0523, 0526–0529, 0982, 0983</p> |

2018 Behavioral Health Guide

| | Appropriate Care and Associated Codes to Bill | | | |
|---|--|---|-------------|------------|
| | CPT Codes | ICD-10 Codes | HCPCS Codes | UB Revenue |
| Follow-up after Emergency Department visits for Mental Illness (FUM) (Continued from page 10) | FUH Visits Group 2 Value Set: 99221–99223, 99231–99233, 99238, 99239, 99251–99255 FUH POS Group 2 Value Set: 52, 53 | F53, F59, F60.0, F60.1, F60.2, F60.3, F60.4, F60.5, F60.6, F60.7, F60.81, F60.89, F60.9 F63.0, F63.1, F63.2, F63.3, F63.81, F63.89, F63.9, F64.0, F64.1, F64.2, F64.8, F64.9, F65.0, F65.1, F65.2, F65.3, F65.4, F65.50, F65.51, F65.52, F65.89, F65.9, F66, F68.10–68.13, F68.8, F69, F80.0, F80.1, F80.2, F80.4, F80.81, F80.82, F80.89, F80.9, F81.0, F81.2, F81.81, F81.89, F81.9, F82, F84.0–F84.3, F84.5, F84.8, F84.9, F88, F89, F90.0, F90.1, F90.2, F90.8, F90.9, F94.0, F94.1, F94.2, F94.8, F94.9, F95.0, F95.1, F95.2, F95.8, F95.9, F98.0, F98.1, F98.21, F98.29, F98.3, F98.4, F98.5, F98.8, F98.9, F99 Mental Health Illness Codes: (To meet this measure the member must have had a ED visit with a principal diagnosis of mental illness and was 6 years or older on the date of service.) F20.0–F20.3, F20.5, F20.81, F20.89, F20.9, F21–24, F25.0, F25.1, F25.8, F25.9, F30.10–F30.13, F30.2–F30.4, F30.8, F30.9, F31.0, F31.10–F31.13, F31.2, F31.30–F31.32, F31.4, F31.5, F31.60–F31.64, F31.70–F31.78, F31.81, F31.89, F31.9, F32.0–F32.5, F32.8, F32.81, F32.89, F32.9, F33.0–F33.3, F33.40–F33.42, F33.8, F33.9, F34.0, F43.1, F34.8, F34.81, F34.89, F34.9, F42, F42.2–F42.4, F421.8, F42.9, F43.0, F43.10–F43.12, F43.20–F43.25, F43.29, F43.8, F43.9, F44.89, F53, F60.0–F60.7, F60.81, F60.89, F60.9, F63.0–F63.3, F63.81, F63.89, F63.9, F68.10–F68.13, F68.8, F84.0, F84.2, F84.3, F84.5, F84.8, F84.9, F90.1, F90.2, F90.8, F90.9, F91.0–F91.3, F91.8, F91.9, F93.0, F93.8, F93.9, F94.0–F94.2, F94.8, F94.9 | | |

2018 Behavioral Health Guide

| | Appropriate Care and Associated Codes to Bill | | | |
|---|---|---|--|--|
| | CPT Codes | ICD-10 Codes | HCPCS Codes | UB Revenue |
| <p>Initiation and Engagement of Alcohol and Other Drug Abuse or Dependence (AOD) Treatment (IET)</p> <p>The percentage of adolescent and adult members with a new episode of alcohol or other drug (AOD) abuse or dependence who received the following:</p> <p>Initiation of AOD Treatment:</p> <p>The percentage of members who initiate treatment through an inpatient AOD admission, outpatient visit, intensive outpatient encounter or partial hospitalization, telehealth or medication assisted treatment (MAT) within 14 days of the diagnosis.</p> <p>OR</p> <p>Engagement of AOD Treatment:</p> <p>The percentage of members who initiated treatment and who had two or more additional AOD services or MAT within 34 days of the initiation visit.</p> <p>Definition: IESD means Index Episode Start Date. For an outpatient, intensive outpatient, partial hospitalization, telehealth, detoxification or ED visit (not resulting in an inpatient stay), the IESD is the date of service.</p> <p>For an inpatient stay, the IESD is the date of discharge.</p> <p>Note: An ED visit that results in an inpatient stay does not require a AOD diagnosis for the stay when identifying the IESD.</p> | <p>ED Codes: 99281–99285</p> <p>Stand Alone Visit Codes need to be with ICD-10 Codes for both initiation or engagement of AOD treatment:</p> <p>IET Stand Alone Visits Value Sets with Alcohol Abuse and Dependence, Opioid Abuse and Dependence or Other Drug Abuse and Dependence with or without Telehealth Modifier Value Set. 98960–98962, 99078, 99201–99205, 99211–99220, 99241–99245, 99341–99345, 99347–99350, 99384–99387, 99394–99397, 99401–99404, 99408, 99409, 99411, 99412, 99510</p> <p>Telehealth Modifier Value Set Codes: 95, or GT</p> <p>IET visits Group 1 Value Set with IET POS Group 1 Value Set and Alcohol Abuse and Dependence, Opioid Abuse and Dependence or Other Drug Abuse and Dependence with or without telehealth modifier.</p> <p>IET Visit Group 1 Value Set: 90791, 90792, 90832–90834, 90836–90840, 90845, 90847, 90849, 90853, 90857, 90876</p> <p>IET POS Group 1 Value Set: 03, 05, 07, 09, 11–20, 22, 33, 49, 50, 52, 53, 57, 71, 72</p> | <p>Alcohol and Other Drug Dependence Diagnosis: F10.10, F10.120, F10.121, F10.129, F10.14, F10.150, F10.151, F10.159, F10.180, F10.181, F10.182, F10.188, F10.19, F10.20, F10.220, F10.221, F10.229–F10.232, F10.239, F10.24, F10.250, F10.251, F10.259, F10.26, F10.27, F10.280–F10.282, F10.288, F10.29,</p> <p>Opioid Abuse and Dependence: F11.10, F11.120, F11.121, F11.122, F11.129, F11.14, F11.150, F11.151, F11.159, F11.181, F11.182, F11.188, F11.19, F11.20, F11.220, F11.221, F11.222, F11.229, F11.23, F11.24, F11.250, F11.251, F11.259, F11.281, F11.282, F11.288, F11.29</p> <p>Other Drug Abuse and Dependence Codes: F12.10, F12.120, F12.121, F12.122, F12.129, F12.150, F12.151, F12.159, F12.180, F12.188, F12.19, F12.20, F12.220, F12.221, F12.222, F12.229, F12.250, F12.251, F12.259, F12.280, F12.288, F12.29, F13.10, F13.120, F13.121, F13.129, F13.14, F13.150, F13.151, F13.159, F13.180, F13.181, F13.182, F13.188, F13.19, F13.20, F13.220, F13.221, F13.229, F13.230, F13.231, F13.232, F13.239, F13.24, F13.250, F13.251, F13.259, F13.26, F13.27, F13.280, F13.281, F13.282, F13.288, F13.29, F14.10, F14.120, F14.121, F14.122, F14.129, F14.14, F14.150, F14.151, F14.159, F14.180, F14.181, F14.182, F14.188, F14.19, F14.20, F14.220, F14.221, F14.222, F14.229, F14.23, F14.24, F14.250, F14.251, F14.259, F14.280, F14.281, F14.282, F14.288, F14.29, F15.10, F15.120, F15.121, F15.122, F15.129,</p> <p><i>(Additional ICD-10 codes on page 13)</i></p> | <p>Use the following IET Stand Alone Visits HCPCS codes with the appropriate ICD-10 codes: G0155, G0176, G0177, G0396, G0397, G0409, G0410, G0411, G0443, G0463, H0001, H0002, H0004, H0005, H0007, H0015, H0016, H0022, H0031, H0034–H0037, H0039, H0040, H0047, H2000, H2000, H2010, H2011, H2013–H2020, H2035, H2036, M0064, S0201, S9480, S9484, S9485, T1006, T1012, T1015</p> <p>If the IESD was for Alcohol Abuse or Dependence use a Alcohol Abuse and Dependence Value Set with the HCPCS for MAT dispensing event or a claim for MAT (Medication Assisted Treatment Value Set): H0020, H0033, J0571–J0575, J2315, S0109</p> <p>Note: Use the MAT value set with diagnosis of Opioid Abuse or dependence if appropriate.</p> | <p>For Acute or Non-Acute Inpatient Admissions use Inpatient Stay alone Value Sets with one of the following ICD-10 value sets: Alcohol Abuse and Dependence, Opioid Abuse and Dependence or Other Drug Abuse and Dependence.</p> <p>Inpatient Codes: 0100, 0101, 0110–0114, 0116–0124, 0126–0134, 0136–0144, 0146–0154, 0156–0160, 0164, 0167, 0169–0174, 0179, 0190–0194, 0199, 0200–0204, 0206–0214, 0219, 1000–1002</p> <p>IET Stand Alone Visits (UBREV): 0510, 0513, 0515–0517, 0519–0523, 0526–0529, 0900, 0902–0907, 0911–0917, 0919, 0944, 0945, 0982, 0983</p> |

2018 Behavioral Health Guide

| Appropriate Care and Associated Codes to Bill | | | | |
|--|--|--|-------------|------------|
| | CPT Codes | ICD-10 Codes | HCPCS Codes | UB Revenue |
| Initiation and Engagement of Alcohol and Other Drug Abuse or Dependence (AOD) Treatment (IET) (Continued from page 12) | <p>IET Visit Group 2 Value Set with IET POS Group 2 Value Set and Alcohol Abuse and Dependence, Opioid Abuse and Dependence or Other Drug Abuse and Dependence Codes with or without telehealth Modifier. A telephone Visit Value Set can also be used with the Alcohol Abuse and Dependence, Opioid Abuse and Dependence or Other Drug Abuse and Dependence.</p> <p>IET Visit Group 2 Value Set: 99221-99223, 99231-99233, 99238, 99239, 99251-99255</p> <p>IET POS Group 2 Value Set: 52, 53</p> <p>Telephone Visit Value Set: 98966-98968, 99441-99443</p> <p>Engagement of AOD Treatment: If the member is compliant for the Initiation of AOD Treatment and two or more inpatient admissions, outpatient visits, telehealth, intensive outpatient encounters or partial hospitalization with a diagnosis matching the IESD diagnosis on the day after the initiation encounter through 34 total days then the member can count for the numerator of the Engagement of AOD Treatment.</p> <p>Note: If multiple engagements occur on the same day they can be counted if with a different provider for each engagement. Follow the same coding sequence outline in the Initiation of AOD Treatment for Engagement of AOD Treatment.</p> <p>Online Assessment Value Set with Alcohol Abuse and Dependence, Opioid Abuse and Dependence or Other Drug Abuse and Dependence: 98969, 99444</p> | F15.14, F15.150, F15.151, F15.159, F15.180, F15.181, F15.182, F15.188, F15.19, F15.20, F15.220, F15.221, F15.222, F15.229, F15.23, F15.24, F15.250, F15.251, F15.259, F15.280, F15.281, F15.282, F15.288, F15.29, F16.10, F16.210, F16.121, F16.122, F16.129, F16.14, F16.150, F16.151, F16.159, F16.80, F16.183, F16.188, F16.19, F16.20, F16.220, F16.221, F16.229, F16.24, F16.250, F16.251, F16.259, F16.280, F16.283, F16.288, F16.29, F18.10, F18.120, F18.121, F18.129, F18.14, F18.150, F18.151, F18.159, F18.17, F18.180, F18.188, F18.19, F18.20, F18.220, F18.221, F18.229, F18.24, F18.250, F18.251, F18.259, F18.27, F18.280, F18.288, F18.29, F19.10, F19.120, F19.121, F19.122, F19.129, F19.14, F19.150, F19.151, F19.159, F19.16, F19.17, F19.180, F19.181, F19.182, F19.188, F19.19, F19.20, F19.220, F19.221, F19.222, F19.229, F19.230, F19.231, F19.232, F19.239, F19.24, F19.250, F19.251, F19.259, F19.26, F19.27, F19.280, F19.281, F19.282, F19.288, F19.29 | | |

2018 Behavioral Health Guide

| | Appropriate Care and Associated Codes to Bill | | | |
|--|--|--|---|--|
| | CPT Codes | ICD-10 Codes | HCPCS Codes | UB Revenue |
| <p>Adherence to Antipsychotic Medications for Individuals with Schizophrenia (SAA)</p> <p>Members who are 19–64 years of age during the measurement year with Schizophrenia who were dispensed and remained on an antipsychotic medication for at least 80% of the treatment period.</p> <p>Identify members with schizophrenia as those who met at least one of the following:</p> <p>At least one acute inpatient encounter with any diagnosis of schizophrenia. Or at least two visits in an outpatient, intensive outpatient, partial hospitalization, ED or non-acute inpatient setting, on different dates of service, with any diagnosis of schizophrenia.</p> | <p>BH Acute Inpatient Value Set with BH Acute Inpatient POS Value Set with Schizophrenia Value Set:</p> <p>90791, 90792, 90832–90834, 90836–90840, 90845, 90847, 90849, 90853, 90867–90870, 90875, 90876, 99221–99223, 99231–99233, 99238, 99239, 99251–99255, 99291</p> <p>BH Acute Inpatient POS Value Set:</p> <p>21, 51</p> <p>BH Stand Alone Outpatient/PH/IOP Value Set Codes:</p> <p>98960–98962, 99078, 99201–99205, 99211–99215, 99217–99220, 99241–99245, 99341–99345, 99347–99350, 99384–99387, 99394–99397, 99401–99404, 99411, 99412, 99510</p> <p>Use the BH Outpatient /PH/IOP Value Set with the BH Outpatient /PH/IOP POS Value Set with the ICD-10 Schizophrenia Value Set:</p> <p>90791, 90792, 90832–90834, 90836–90840, 90845, 90847, 90849, 90853, 90867–90870, 90875, 90876, 99221–99223, 99231–99233, 99238, 99239, 99251–99255, 99291</p> <p>BH Outpatient /PH/ IOP POS Value Set:</p> <p>03, 05, 07, 09, 11–20, 22, 24, 33, 49, 50, 52, 53, 71, 72</p> <p>Use the CPT ED Value Set with the ICD-10 Schizophrenia Value Set:</p> <p>99281–99285</p> <p><i>(Additional CPT codes on page 15)</i></p> | <p>Schizophrenia codes only, exclude dementia codes:</p> <p>F20.0–F20.3, F20.5, F20.81, F20.89, F20.9, F25.0, F25.1, F25.8, F25.9</p> | <p>Use HCPCS codes for the BH Stand Alone Outpatient/PH/IOP Value Set with the Schizophrenia ICD-10 codes:</p> <p>G0155, G0176, G0177, G0409–G0411, G0463, H0002, H0004, H0031, H0034–HH0037, H0039, H0040, H2000, H2001, H2010–HH2020, M0064, S0201, S9480, S9484, S9485, T1015</p> <p>Use these HCPCS codes for the BH Stand Alone Non-acute Inpatient Value Set with ICD-10 Schizophrenia Value Set:</p> <p>H0017–H0019, T2048</p> <p>Long Acting Injections 14 day supply Value Set:</p> <p>J2794</p> <p>Long Acting Injections 28 Day Supply Value Set:</p> <p>J0401, J1631, J2358, J2426, J2680</p> | <p>Use UB Rev codes for Behavioral Health (BH) Stand Alone Acute Inpatient Value Set with Schizophrenia Value Set under ICD-10 column.</p> <p>0100, 0101, 0110, 0111–0114, 0119–0124, 0129–0134, 0139–0144, 0149–0154, 0160, 0164, 0167, 0169, 0200–0204, 0206–0214, 0219, 0720–0724, 0729, 0987</p> <p>Use UB Rev ED Value Set with the ICD-10 Schizophrenia Value Set:</p> <p>0450–0452, 0456, 0459, 0981</p> <p>UB Rev BH Stand Alone Non-acute Inpatient Value Set with the ICD-10 Schizophrenia Value Set:</p> <p>0118, 0128, 0138, 0148, 0158, 0190–0194, 0199, 0524, 0525, 0550–0552, 0559–0663, 1000, 1001, 1003–1005</p> |

2018 Behavioral Health Guide

| | Appropriate Care and Associated Codes to Bill | | | |
|--|---|--------------|-------------|------------|
| | CPT Codes | ICD-10 Codes | HCPCS Codes | UB Revenue |
| Adherence to Antipsychotic Medications for Individuals with Schizophrenia (SAA) (Continued from page 14) | <p>Use the BH ED Value Set with the ED POS Value Set with the ICD-10 Schizophrenia Value Set: 90791, 90792, 90832–90834, 90836–90840, 90845, 90847, 90849, 90853, 90867–90870, 90875, 90876, 99291</p> <p>ED POS Value Set: 23</p> <p>Use the BH Stand Alone Non-acute Inpatient CPT Value Set with the ICD-10 Schizophrenia Value Set: 99304–99310, 99315, 99319, 99318, 99324–99328, 99334–99337</p> <p>Use BH Non-acute Inpatient CPT Value Set with BH Non-acute Inpatient POS Value Set with the ICD-10 Schizophrenia Value Set: 90791, 90792, 90832–90834, 90845, 90847, 90849, 90853, 90867–90870, 90875, 90876, 99291</p> <p>BH Non-acute Inpatient POS value Set: 31, 32, 56</p> | | | |

2018 Behavioral Health Guide

| | Appropriate Care and Associated Codes to Bill | | | |
|--|---|---|--|--|
| | CPT Codes | ICD-10 Codes | HCPCS Codes | UB Revenue |
| <p>Cardiovascular Monitoring for People with Cardiovascular Disease and Schizophrenia (SMC)</p> <p>Members 18–64 years of age with Schizophrenia and Cardiovascular Disease, who had an LDL-C test during the measurement year.</p> <p>For the purpose of this measure, cardiovascular disease is defined either as a diagnosis of IVD (Ischemic Vascular Disease) or Inpatient treatment for AMI, CABG, or PCI.</p> | <p>LDL-C Screening: 80061, 83700, 83701, 83704, 83721</p> <p>LDL-C Tests for LOINC: 12773–8, 13457–7, 18261–8, 18262–6, 2089–1, 49132–4, 55440–2</p> <p>LDL-C Tests for CPT-CAT-II codes: 3048F, 3049F, 3050F</p> <p>BH Stand Alone Outpatient/PH/IOP Codes with ICD-10 Schizophrenia Codes: 98960–98962, 99078, 99201–99205, 99211–99215, 99217–99220, 99241–99245, 99341–99345, 99347–99350, 99384–99387, 99394–99397, 99401–99404, 99411, 99412, 99510</p> <p>BH Outpatient/PH/IOP Codes used with BH Outpatient/PH/IOP/POS Codes and with a Schizophrenia ICD-10 Code: 90791, 90792, 90832, 90833, 90834, 90836–90840, 90845, 90847, 90849, 90853, 90867, 90868, 90869, 90870, 90875, 90876, 99221–99223, 99231–99233, 99238, 99239, 99251–99255, 99291</p> <p>Add the following BH Outpatient/PH/IOP POS Value Set codes: 03, 05, 07, 09, 11–20, 22, 24, 33, 49–53, 71, 72</p> <p><i>(Additional CPT codes on page 17)</i></p> | <p>Schizophrenia Codes: F20.0–F20.3, F20.5, F20.81, F20.89, F20.9, F25.0, F25.1, F25.8, F25.9</p> <p>Note: members are identified as having cardiovascular disease in two ways: by event or by diagnosis.</p> <p>For Event: Code AMI Value Set with the Inpatient Stay Value Set: AMI Codes are: 121.01, 121.02, 121.09, 121.11, 121.19, 121.21, 121.29, 121.3, 121.4</p> <p>For Diagnosis: Ischemic Vascular Disease (IVD) value set: I20.0, I20.8, I20.9, I24.0, I24.8, I24.9, I25.10, I25.110, I25.111, I25.118, I25.119, I25.5, I25.6, I25.700, I25.701, I25.708–I25.711, I25.718, I25.71, I25.720, I25.721, I25.728, I25.729, I25.730, I25.731, I25.738, I25.739, I25.750, I25.751, I25.758, I25.759, I25.760, I25.761, I25.768, I25.769, I25.790, I25.791, I25.798, I25.799, I25.810, I25.811, I25.812, I25.82, I25.83, I25.84, I25.89, I25.9, I63.20, I63.211, I63.212, I63.219, I63.22, I63.231, I63.232, I63.239, I63.29, I63.50, I63.511, I63.512, I63.513, I63.519, I63.521, I63.522, I63.523, I63.529, I63.531, I63.532, I63.533, I63.539, I63.541, I63.542, I63.543, I63.549, I63.59, I65.01, I65.02, I65.03, I65.09, I65.1, I65.21, I65.22, I65.23, I65.29, I65.8, I65.9, I66.01–I66.03, I66.09, I66.11, I66.12, I66.13, I66.19, I66.21, I66.22, I66.23, I66.29, I66.3, I66.8, I66.9, I67.2, I70.1, I70.201–I70.203, I70.208, I70.209, I70.211–I70.213, I70.218, I70.219, I70.221, I70.222, I70.223, I70.228, I70.229,</p> <p><i>(Additional ICD-10 codes on page 17)</i></p> | <p>Use the HCPCS codes with the BH Stand Alone Outpatient/PH/IOP codes with the schizophrenia codes under ICD-10: G0155, G0176, G0177, G0409–G0411, G0463, H0002, H0004, H0031, H0034, H0035, H0036, H0037, H0039, H0040, H2000, H2001, H2010–H2020, M0064, S0201, S9480, S9484, S9485, T1015</p> <p>BH Stand Alone Nonacute Inpatient Value Set with Schizophrenia Value Set: H0017–H0019, T2048</p> <p>HCPCS CABG Value Set Codes: S2205–S2209</p> <p>HCPCS PCI Value Set Codes: C9600, C9602, C9604, C9606, C9607</p> <p>HCPCS Outpatient Value Set Codes use with the IVD ICD-10 Value Set Codes: G0402, G0438, G0439, G0463, T1015</p> | <p>At least one acute inpatient encounter with any diagnosis of schizophrenia using the BH stand alone acute inpatient value set with the schizophrenia value set: 0100, 0101, 0110–0114, 0119–0124, 0129–0134, 0139–0144, 0149–0154, 0159, 0160, 0164, 0167, 0169–0204, 0206–0214, 0219, 0720–0724, 0729, 0987</p> <p>Must have at least two visits in an outpatient, intensive outpatient, partial hospitalization, ED or nonacute inpatient setting on different dates use the following codes:</p> <p>BH stand alone outpatient/PH/IOP Value Set. Use with Schizophrenia Value Set Codes: 0510, 0513, 0516, 0517, 0519–0523, 0526–0259, 0900–0905, 0907, 0911–0917, 0919, 0982, 0983</p> <p>Code BH Stand Alone Nonacute Inpatient Value Set with Schizophrenia ICD-10 Value Set: 0118, 0128, 0138, 0148, 0158, 0190–0194, 0199, 0524, 0525, 0550–0552, 0559–0663, 0669, 1000–1005,</p> <p>Inpatient Stay Value Set Codes. Use with AMI Value Set and IVD Value Set: 0100, 0101, 0110–0124, 0126–0134, 0136–0144, 0146–0154, 0156–0160, 0164, 0167, 0169–0174, 0179, 0190–0194, 0199–0204, 0206–0214, 0219, 1000, 1001, 1002</p> <p><i>(Additional UB Revenue codes on page 17)</i></p> |

2018 Behavioral Health Guide

| Appropriate Care and Associated Codes to Bill | | | | |
|---|--|--|-------------|---|
| | CPT Codes | ICD-10 Codes | HCPCS Codes | UB Revenue |
| Cardiovascular Monitoring for People with Cardiovascular Disease and Schizophrenia (SMC) (Continued from page 16) | <p>At least one acute inpatient encounter with any diagnosis of schizophrenia using BH acute inpatient value set with BH acute inpatient POPS value set with schizophrenia value set: 90791, 90792, 90832–90834, 90836–90840, 90845, 90847, 90849, 90853, 90867–90870, 90875, 90876, 99221–99223, 99231–99233, 99238, 99239, 99251–99255, 99291</p> <p>BH acute inpatient POS value set: 21, 51</p> <p>ED Value Set with Schizophrenia ICD-10 Value Set Codes: 0450–0452, 0456, 0459, 0981, 99281–99285</p> <p>Code the BH ED Value set with ED POS Value Set with Schizophrenia ICD-10 Value Set: 90791, 90792, 90832–90834, 90836–90840, 90845, 90847, 90849, 90853, 90867, 90869, 90870, 90875, 90876, 99291</p> <p>ED POS Value Set: 23</p> <p>BH Stand Alone Nonacute Inpatient Value Set with Schizophrenia ICD-10 Value Set: 99304–99310, 99315, 99316, 99318, 99324–99328, 99334–99337</p> <p><i>(Additional CPT codes on page 18)</i></p> | I70.231–I70235, I70.238, I70.239, I70.241–I70.245, I70.248, I70.249, I70.25, I70.261–I70.263, I70.268, I70.269, I70.291–I70.293, I70.298, I70.299, I70.301, I70.302, I70.303, I70.308–I70.313, I70.318, I70.319, I70.321, I70.322, I70.323, I70.328, I70.329, I70.331, I70.332, I70.333, I70.334, I70.335, I70.338, I70.339, I70.341–I70.345, I70.348, I70.349, I70.35, I70.361, I70.362, I70.363, I70.368, I70.369, I70.391, I70.392, I70.393, I70.398, I70.399, I70.401, I70.402, I70.403, I70.408, I70.409, I70.411, I70.412, I70.413, I70.418, I70.419, I70.421, I70.422, I70.423, I70.428, I70.429, I70.431–I70435, I70.438, I70.439, I70.441, I70.442, I70.443, I70.444, I70.445, I70.448, I70.449, I70.45, I70.461, I70.462, I70.463, I70.468, I70.469, I70.491, I70.492, I70.493, I70.498, I70.499, I70.501–I70.503, I70.508, I70.509, I70.511, I70.512, I70.513, I70.518, I70.519, I70.521–I70.523, I70.528, I70.529, I70.531–I70.535, I70.538, I70.539, I70.541–I70.545, I70.548, I70.549, I70.55, I70.561, I70.562, I70.563, I70.568, I70.569, I70.591, I70.592, I70.593, I70.598, I70.599, I70.601, I70.602, I70.603, I70.608, I70.609, I70.611–I70613, I70.618, I70.619, I70.621–I70.623, I70.628, I70.629, I70.631–I70.635, I70.638, I70.639, I70.641–I70.645, I70.648, I70.649, I70.65, I70.661, I70.662, I70.663, I70.668, I70.669, I70.691, I70.692, I70.693, I70.698, I70.699, I70.701, I70.702, I70.703, I70.708, I70.709, I70.711–I70.713, I70.718, I70.719, I70.721, I70.722, I70.723, I70.728, I70.729, I70.731–I70.735, I70.738, I70.739, I70.741–I70745, I70.748, I70.749, I70.75, I70.761–I70.763, I70.768, I70.769, I70.791–I70.793, I70.798, I70.799, I70.92, I75.011, I75.012, I75.013, I75.019, I75.021, I75.022, I75.023, I75.029, I75.81, I75.89, T82.855A, T82.855D, T82.855S, T82.856A, T82.856D, T82.856S <i>(Additional ICD-10 codes on page 18)</i> | | <p>Outpatient Value Set Codes use with the IVD Value Set: 0510–0517, 0519–0523, 0526–0529, 0982, 0983</p> <p>Acute Inpatient Value Set Codes to be used with IVD Value Set codes under the ICD-10 column: 0100, 0101, 0110–0114, 0119–0124, 0129–0134, 0139–0144, 0149–0154, 0159–0160, 0164, 0167, 0169, 0200–0204, 0206–0214, 0219, 0720–0724, 0729, 0987</p> |

2018 Behavioral Health Guide

| | Appropriate Care and Associated Codes to Bill | | | |
|---|--|--|-------------|------------|
| | CPT Codes | ICD-10 Codes | HCPCS Codes | UB Revenue |
| Cardiovascular Monitoring for People with Cardiovascular Disease and Schizophrenia (SMC) (Continued from page 17) | BH Nonacute Inpatient Value Set with BH Nonacute Inpatient POS Value Set with Schizophrenia ICD-10 Value Set: 90791, 90792, 90832–90834, 90836–90840, 90845, 90847, 90849, 90853, 90867–90870, 90875, 90876, 99291 BH Nonacute Inpatient POS Value Set: 31, 32, 56 Acute Inpatient Value Set CPT codes to be used with the IVD ICD-10 Value Set Codes: 99221–99223, 99231–99233, 99238, 99239, 99251–99255, 99291 CPT CABG Value Set Codes: 33510–33514, 33516–33519, 33521–33523, 33533–33536 CPT PCI Value Set Codes: 92920, 92924, 92928, 92933, 92937, 92941, 92943, 92980, 92982, 92995 Outpatient Value Set Codes use with the IVD Value Set: 99201–99205, 99211–99215, 99241–99245, 99341–99345, 99347–99350, 99381–99387, 99392–99397, 99401–99404, 99411, 99412, 99456 | CABG Value Set: 0210083, 0210088, 0210089, 021008C, 021008F, 021008W, 0210093, 0210098, 0210099, 021009C, 021009F, 021009W, 02100A3, 02100A8, 02100A9, 02100AC, 02100AF, 02100AW, 02100J3, 02100J8, 02100J9, 02100JC, 02100JF, 02100JW, 02100K3, 02100K8, 02100K9, 02100KC, 02100KF, 02100KW, 02100Z3, 02100Z8, 02100Z9, 02100ZC, 02100ZF, 0211083, 0211088, 0211089, 021108C, 021108F, 021108W, 0211093, 0211098, 0211099, 021109C, 021109F, 021109F, 021109W, 02110A3, 02110A8, 02110A9, 02110AC, 02110AF, 02110AW, 02110J3, 02110J8, 02110J9, 02110JC, 02110JF, 02110JW, 02110K3, 02110K8, 02110K9, 02110KC, 02110KF, 02110KW, 02110Z3, 02110Z8, 02110Z9, 02110ZC, 02110ZF, 0212083, 0212088, 0212089, 021208C, 021208F, 021208W, 0212093, 0212098, 0212099, 021209C, 021209C, 021209F, 021209W, 02120A3, 02120A8, 02120A9, 02120AC, 02120AF, 02120AW, 02120J3, 02120J8, 02120J9, 02120JC, 02120JF, 02120JW, 02120K3, 02120K8, 02120K9, 02120KC, 02120KF, 02120KW, 02120Z3, 02120Z8, 02120Z8, 02120Z9, 02120ZC, 02120ZF, 0213083, 0213088, 0213089, 021308C, 021308F, 021308W, 0213093, 0213098, 0213099, 021309C, 021309F, 021309W, 02130A3, 02130A8, 02130A9, 02130AC, 02130AF, 02130AW, 02130J3, 02130J8, 02130J9, 02130JC, 02130JF, 02130JW, 02130K3, 02130K8, 02130K8, 02130K9, 02130KC, 02130KF, 02130KW, 02130Z3, 02130Z8, 02130Z9, 02130ZC, 02130F <i>(Additional ICD-10 codes on page 19)</i> | | |

2018 Behavioral Health Guide

| Appropriate Care and Associated Codes to Bill | | | |
|---|---|-------------|------------|
| CPT Codes | ICD-10 Codes | HCPCS Codes | UB Revenue |
| Cardiovascular Monitoring for People with Cardiovascular Disease and Schizophrenia (SMC) (Continued from page 18) | ICD-10 PCI Value Set: 0270346, 027034Z, 0270356, 027035Z, 0270366, 027036Z, 0270376, 027037Z, 02703D6, 02703DZ, 02703E6, 027023EZ, 02703F6, 02703FZ, 02703G6, 02703T6, 02703TZ, 02703Z6, 02703ZZ, 0270446, 027044Z, 0270456, 027045Z 0270466, 027046Z, 0270476, 027047Z, 02704D6, 02704DZ, 02704E6, 02704EZ, 02704F6, 02704FZ, 02704G6, 02704GZ, 02704T6, 02704TZ, 02704Z6, 02704ZZ, 0271346, 027134Z, 0271356, 027135Z, 0271366, 027136Z, 0271376, 0271137Z, 02713D6, 02713DZ, 02713E6, 02713EZ, 02713F6, 0713FZ, 02713G6, 02713GZ, 02713T6, 02713TZ, 02713Z6, 02713ZZ, 0271446, 027144Z, 0271456, 027145Z, 0271466, 027146Z, 0271476, 027147Z, 02714D6, 02714DZ, 02714E6, 02714EZ, 02714F6, 02714FZ, 02714G6, 02714GZ, 02714T6, 02714TZ, 02714Z6, 02714ZZ, 0272346, 027234Z, 0272356, 027235Z, 0272366, 027236Z, 0272376, 027237Z, 02723D6, 02723DZ, 02723E6, 02723EZ, 02723F6, 02723FZ, 02723G6, 02723GZ, 02723T6, 02723TZ, 02723Z6, 02723ZZ, 0272446, 027244Z, 0272456, 027245Z, 0272466, 027246Z, 0272476, 027247Z, 02724D6, 02724DZ, 02724E6, 02724EZ, 02724F6, 02724FZ, 02724G6, 02724GZ, 02724T6, 02724TZ, 02724Z6, 02724ZZ, 0273346, 027334Z, 0273356, 027335Z, 0273366, 027336Z, 0273376, 027337Z, 02733D6, 02733DZ, 02733E6, 02733EZ, 02733F6, 02733FZ, 02733G6, 02733GZ, 02733T6, 02733TZ, 02733Z6, 02733ZZ, 0273446, 027344Z, 0273456, 027345Z, 0273466, 027346Z, 0273476, 027347Z, 02734D6, 02734DZ, 02734E6, 02734EZ, 02734F6, 02734FZ, 02734G6, 02734GZ, 02734T6, 02734TZ, 02734Z6, 02734ZZ | | |

2018 Behavioral Health Guide

| Appropriate Care and Associated Codes to Bill | | | | |
|--|---|---|--|---|
| | CPT Codes | ICD-10 Codes | HCPCS Codes | UB Revenue |
| Diabetes Monitoring for People with Diabetes and Schizophrenia (SMD) Members 18–64 years of age with Schizophrenia and Diabetes, who had both an LDL-C test and an HbA1c test during the measurement year. | HbA1c Test: 83036, 83037 LOINC codes for HbA1c Tests: 17856–6, 4548–4, 4549–2 CPT-CAT-II codes for HbA1c Tests: 3044F, 3045F, 3046F LDL-C Tests: 80061, 83700, 83701, 83704, 83721 LDL-C LOINC codes: 12773–8, 13457–7, 18261–8, 18262–6, 49132–4, 55440–2 LDL-C CPT-CAT-II Codes: 3048F, 3049F, 3050F Code at least one BH Acute Inpatient Value Set with BH Acute Inpatient POS Value Set and with the ICD-10 Schizophrenia Value Set: 90791, 90792, 90832–90834, 90836–90840, 90845, 90847, 90849, 90853, 90867, 90868–90870, 90875, 90876, 99221–99223, 99231–99233, 99238, 99239, 99251–99255, 99291 BH Stand Alone Outpatient/PH/IOP Codes with the Schizophrenia ICD-10 Codes: 98960–98962, 99078, 99201–99205, 99211–99215, 99217–99220, 99241–99245, 99341–99345, 99347–99350, 99384–99387, 99394–99397, 99401–99404, 99411, 99412, 99510 <i>(Additional CPT codes on page 21)</i> | Schizophrenia codes only: F20.0–F20.3, F20.5, F20.81, F20.89, F20.9, F25.0, F25.1, F25.8, F25.9 Diabetes: E10.10, E10.11, E10.21, E10.22, E10.29, E10.311, E10.319, E10.321, E10.3211, E10.3212, E10.3213, E10.3219, E10.329, E10.3291–E10.3293, E10.3299, E10.331, E10.311–E10.3313, E10.3319, E10.339, E10.3391–E10.3393, E10.3399, E10.341, E10.3411–E10.3413, E10.3419, E10.349, E10.3491–E10.3493, E10.3499, E10.351, E10.3511–E10.3513, E10.3519, E10.3521–E10.3523, E10.3529, E10.3531–E10.3533, E10.3539, E10.3541–E10.3543, E10.3549, E10.3551–E10.3553, E10.3559, E10.359, E10.3591–E10.3593, E10.3599, E10.36, E10.37X1–E10.37X3, E10.37X9, E10.39, E10.40, E10.41, E10.42, E10.43, E10.44, E10.49, E10.51, E10.52, E10.59, E10.610, E10.618, E10.620, E10.621, E10.622, E10.628, E10.630, E10.638, E10.641, E10.649, E10.65, E10.69, E10.8, E10.9, E11.00, E11.01, E11.21, E11.22, E11.29, E11.311, E11.319, E11.321, E11.3211–E11.3213, E11.3219, E11.329, E11.3291–E11.3293, E11.3299, E11.331, E11.3311–E11.3313, E11.3319, E11.339, E11.3391–E11.3393, E11.3399, E11.341, E11.3411–E11.3413, E11.3419, E11.349, E11.3491, E11.3492, E11.3493, E11.3499, E11.351, E11.3511–E11.3513, E11.3519–E11.3523, E11.3529, E11.3531–E11.3533, E11.3539, E11.3541–E11.3543, E11.3549, E11.3551–E11.3553, E11.3559, E11.359, E11.3591, E11.3592, E11.3593, E11.3599, E11.36, E11.37X1–E11.37X3, E11.37X9, <i>(Additional ICD-10 codes on page 21)</i> | Use HCPCS BH Stand Alone Outpatient /PH/IOP Value Set Codes with Schizophrenia ICD-10 Value Set Codes: G0155, G0176, G0177, G0409–G0411, G0463, H0002, H0004, H0031, H0034, H0035, H0036, H0037, H0039, H0040, H2000, H2001, H2010–H2020, M0064, S0201, S9480, S9484, S9485, T1015 BH Stand Alone Nonacute Inpatient Value Set with Schizophrenia Value Set: H0017–H0019, T2048 Use the HCPCS Outpatient Value Set Codes with the Diabetes ICD-10 Value Set Codes: G0402, G0438, G0439, G0463, T1015 | BH Stand Alone Acute Inpatient Value Set with the Schizophrenia ICD-10 Value Set Codes: 0100, 0101, 0110–0114, 0119–0124, 0129–0134, 0139–0144, 0149–0154, 0159–0160, 0164, 0167, 0169, 0200–0204, 0206–02124, 0219, 0720–0724, 0729, 0987 Code UB Revenue BH Stand Alone Outpatient /PH/IOP Value Set with Schizophrenia ICD-10 Value Set Codes: 0510, 0513, 0516, 0517, 0519–0523, 0526–0529, 0900–0905, 0907, 0911–0917, 0919, 0982, 0983 ED UB Revenue Value Set Codes with the Schizophrenia ICD-10 Value Set Codes or the UB Revenue ED Value Set Codes with the Diabetes ICD-10 Codes: 0450–0452, 0456, 0459, 0981 Code BH Stand Alone Nonacute Inpatient Value Set with Schizophrenia ICD-10 Value Set: 0118, 0128, 0138, 0148, 0158, 0190–0194, 0199, 0524, 0525, 0550–0552, 0559–0663, 0669, 1000–1005 UB Revenue Outpatient Value Set with the Diabetes ICD-10 Value Set: 0510–0517, 0519–0523, 0526–0529, 0982, 0983 <i>(Additional UB Revenue codes on page 21)</i> |

2018 Behavioral Health Guide

| | Appropriate Care and Associated Codes to Bill | | | |
|--|--|--|-------------|--|
| | CPT Codes | ICD-10 Codes | HCPCS Codes | UB Revenue |
| <p>Diabetes Monitoring for People with Diabetes and Schizophrenia (SMD)</p> <p>Members 18–64 years of age with Schizophrenia and Diabetes, who had both an LDL-C test and an HbA1c test during the measurement year.</p> <p>(Continued from page 20)</p> | <p>BH Outpatient/PH/IOP Codes used with BH Outpatient/PH/IOP/POS Codes and with a Schizophrenia ICD-10 Code:</p> <p>90791, 90792, 90832, 90833, 90834, 90836–90840, 90845, 90847, 90849, 90853, 90867, 90868, 90869, 90870, 90875, 90876, 99221–99223, 99231–99233, 99238, 99239, 99251–99255, 99291</p> <p>BH Outpatient /PH/ IOP POS Value Set:</p> <p>03, 05, 07, 09, 11–20, 24, 33, 49, 50, 52, 53, 71, 72</p> <p>CPT Acute Inpatient Value Set with Diabetes ICD-10 Codes:</p> <p>99221–99223, 99231–99233, 99238, 99239, 99251–99255, 99291</p> <p>ED CPT Value Set Codes with the Schizophrenia ICD-10 Value Set Codes or ED CPT Value Set Codes with the Diabetes ICD-10 Codes:</p> <p>99281–99285</p> <p>Code the BH ED CPT Value Set with ED POS Value Set with the Schizophrenia Value Set Codes:</p> <p>90791, 90792, 90832–90834, 90836–90840, 90845, 90847, 90849, 90853, 90867, 90869, 90870, 90875, 90876, 99291</p> <p>ED POS Value Set:</p> <p>23</p> <p>BH Stand Alone Nonacute Inpatient Value Set with Schizophrenia ICD-10 Value Set:</p> <p>99304–99310, 99315, 99316, 99318, 99324–99328, 99334–99337</p> <p><i>(Additional CPT codes on page 22)</i></p> | <p>E11.39, E11.40, E11.41, E11.42, E11.43, E11.44, E11.49, E11.51, E11.52, E11.59, E11.610, E11.618, E11.620, E11.621, E11.622, E11.628, E11.630, E11.638, E11.641, E11.649, E11.65, E11.69, E11.8, E11.9, E13.00, E13.01, E13.10, E13.11, E13.21, E13.22, E13.29, E13.311, E13.319, E13.321, E13.3211–E13.3213, E13.3219, E13.329, E13.3291–E13.3293, E13.3299, E13.331, E13.3311–E13.3313, E13.3319, E13.339, E13.3391–E13.3393, E13.3399, E13.341, E13.3411–E13.3413, E13.3419, E13.349, E13.3491–E13.3493, E13.3499, E13.351, E13.3511–E13.3513, E13.3519, E13.3521–E13.3523, E13.3529, E13.3531–E13.3533, E13.3539, E13.3541–E13.3543, E13.3549, E13.3551–E13.3553, E13.3559, E13.359, E13.3591, E13.3592, E13.3593, E13.3599, E13.36, E13.37X1–E13.37X3, E13.37X9, E13.39, E13.40, E13.41, E13.42, E13.43, E13.44, E13.49, E13.51, E13.52, E13.59, E13.610, E13.618, E13.620, E13.621, E13.622, E13.628, E13.630, E13.638, E13.641, E13.649, E13.65, E13.69, E13.8, E13.9, O24.011, O24.012, O24.013, O24.019, O24.02, O24.03, O24.111, O24.112, O24.113, O24.119, O24.12, O24.13, O24.311, O24.312, O24.313, O24.319, O24.32, O24.33, O24.811, O24.812, O24.813, O24.819, O24.82, O24.83,</p> | | <p>Nonacute UB Revenue Value Set Codes with the Diabetes ICD-10 Codes:</p> <p>0118, 0128, 0138, 0148, 0158, 0190–0194, 0199, 0524, 0525, 0550–0552, 0559–0663, 0669</p> <p>UB Revenue Acute Inpatient Value Set with the Diabetes ICD-10 Value Set:</p> <p>0100, 0101, 0110–0114, 0119–0124, 0129–0134, 0139–0144, 0149–0154, 0159–0160, 0164, 0167, 0169, 0200–0204, 0206–0214, 0219, 0720–0724, 0729, 0987</p> |

2018 Behavioral Health Guide

| Appropriate Care and Associated Codes to Bill | | | | |
|---|--|--------------|-------------|------------|
| | CPT Codes | ICD-10 Codes | HCPCS Codes | UB Revenue |
| <p>Diabetes Monitoring for People with Diabetes and Schizophrenia (SMD)</p> <p>Members 18–64 years of age with Schizophrenia and Diabetes, who had both an LDL-C test and an HbA1c test during the measurement year.</p> <p>(Continued from page 21)</p> | <p>BH Nonacute Inpatient Value Set with BH Nonacute Inpatient POS Value Set with Schizophrenia ICD-10 Value Set:</p> <p>90791, 90792, 90832–90834, 90836–90840, 90845, 90847, 90849, 90853, 90867–90870, 90875, 90876, 99291</p> <p>Use The CPT Outpatient value Set Codes with the Diabetes ICD-10 Value Set Codes:</p> <p>99201–99205, 99211–99215, 99241–99245, 99341–99345, 99347–99350, 99381–99387, 99392–99397, 99401–99404, 99411, 99412, 99456</p> <p>CPT Observation Value set Codes with the Diabetes ICD-10 Codes:</p> <p>99217–99220</p> <p>CPT nonacute inpatient value set codes with the diabetes ICD-10 code:</p> <p>99304–99310, 99315, 99316, 99318, 99324–99328, 99334–99337</p> | | | |

2018 Behavioral Health Guide

| | Appropriate Care and Associated Codes to Bill | | | |
|---|--|--|---|--|
| | CPT Codes | ICD-10 Codes | HCPCS Codes | UB Revenue |
| <p>Diabetes Screening for People with Schizophrenia or Bipolar Disorder who are using Antipsychotic Medications (SSD)</p> <p>Members 18–64 years of age with Schizophrenia or Bipolar Disorder, who were dispensed an antipsychotic medications and had a diabetes screening test during the measurement year.</p> | <p>Glucose Tests: 80047, 80048, 80050, 80053, 80069, 82947, 82950, 82951</p> <p>LOINC Glucose Tests: 17865–7, 20436–2, 20437–0, 20438–8, 25666–9, 26554–6, 30251–3, 30265–3, 30267–9, 32320–4, 40285–9, 40286–7, 41024–1, 49134–0, 51597–1, 55351–1, 55381–8, 6749–6, 74774–1</p> <p>HbA1c Tests: 83036–83037</p> <p>HbA1c LOINC Codes: 17856–6, 4548–4, 4549–2</p> <p>HbA1c CPT-CAT-II Codes: 3044F, 3045F, 3046F</p> <p>BH Stand Alone Outpatient/PH/IOP codes with the Schizophrenia, or Bipolar or other Bipolar ICD-10 diagnosis codes: 98960–98962, 99078, 99201–99205, 99211–99215, 99217–99220, 99241–99245, 99341–99345, 99347–99350, 99384, 99385, 99386, 99395, 99396, 99401–99404, 99411, 99412, 99510</p> <p><i>(Additional CPT codes on page 24)</i></p> | <p>Schizophrenia codes Only: F20.0–F20.3, F20.5, F20.81, F20.89, F20.9, F25.0, F25.1, F25.8, F25.9</p> <p>Bipolar Disorder: F30.10–F30.13, F30.2–F30.4, F30.8, F30.9, F31.0, F31.10–F31.13, F31.2, F31.30–F31.32, F31.4, F31.5, F31.60–F31.64, F31.70–F31.78</p> <p>Other Bipolar Disorder Value Set Codes: F31.81, F31.89, F31.9</p> | <p>Use HCPCS BH Stand Alone Outpatient /PH/IOP Value Set Codes with Schizophrenia ICD-10 Value Set Codes: G0155, G0176, G0177, G0409–G0411, G0463, H0002, H0004, H0031, H0034, H0035, H0036, H0037, H0039, H0040, H2000, H2001, H2010–H2020, M0064, S0201, S9480, S9484, S9485, T1015</p> <p>BH Stand Alone Nonacute Inpatient Value Set with Schizophrenia, or Bipolar or Other Bipolar ICD-10 Value Set: H0017–H0019, T2048</p> | <p>BH Stand Alone Acute Inpatient Value Set with the Schizophrenia or Bipolar, or Other Bipolar ICD-10 Value Set Codes: 0100, 0101, 0110–0114, 0119–0124, 0129–0134, 0139–0144, 0149–0154, 0159–0160, 0164, 0167, 0169, 0200–0204, 0206–0214, 0219, 0720–0724, 0729, 0987</p> <p>Code UB Revenue BH Stand Alone Outpatient /PH/IOP Value Set with Schizophrenia, or Bipolar, or Other Bipolar ICD-10 Value Set Codes: 0510, 0513, 0516, 0517, 0519–0523, 0526–0529, 0900–0905, 0907, 0911–0917, 0919, 0982, 0983</p> <p>Code BH Stand Alone Nonacute Inpatient Value Set with Schizophrenia, or Bipolar or Other Bipolar ICD-10 Value Set: 0118, 0128, 0138, 0148, 0158, 0190–0194, 0199, 0524, 0525, 0550–0552, 0559–0663, 0669, 1000–1005</p> |

2018 Behavioral Health Guide

| | Appropriate Care and Associated Codes to Bill | | | |
|---|---|--------------|-------------|------------|
| | CPT Codes | ICD-10 Codes | HCPCS Codes | UB Revenue |
| <p>Diabetes Screening for People with Schizophrenia or Bipolar Disorder who are using Antipsychotic Medications (SSD) (Continued from page 23)</p> | <p>BH Outpatient/Ph/IOP Value Set codes use with BH Outpatient / PH/IOP POS Value Set codes with Schizophrenia, or Bipolar or Other Bipolar ICD-10 Diagnosis Codes: 90791, 90792, 90832, 90833, 90834, 90836–90840, 90845, 90847, 90849, 90853, 90867, 90868, 90869, 90870, 90875, 90876, 99221–99223, 99231–99233, 99238, 99239, 99251–99255, 99291</p> <p>BH Outpatient /PH/IOP POS Value Set use with Schizophrenia or Bipolar or Other Bipolar diagnosis ICD-10 Codes: 03, 05, 07, 09, 11–20, 24, 33, 49, 50, 52, 53, 71, 72</p> <p>Code at least one BH Acute Inpatient Value Set with BH Acute Inpatient POS Value Set and with the ICD-10 Schizophrenia, or Bipolar or Other Bipolar Value Set: 90791, 90792, 90832–90834, 90836–90840, 90845, 90847, 90849, 90853, 90867, 90868–90870, 90875, 90876, 99221–99223, 99231–99233, 99238, 99239, 99251–99255, 99291</p> <p>BH Acute Inpatient POS Code: 21, 51</p> <p>Code the BH ED CPT Value Set with ED POS Value Set with the Schizophrenia, or Bipolar or Other Bipolar ICD-10 Value Set Codes: 90791, 90792, 90832–90834, 90836–90840, 90845, 90847, 90849, 90853, 90867, 90869, 90870, 90875, 90876, 99291</p> <p>ED POS Value Set: 23</p> <p><i>(Additional CPT codes on page 25)</i></p> | | | |

2018 Behavioral Health Guide

| Appropriate Care and Associated Codes to Bill | | | | |
|---|--|--------------|-------------|------------|
| | CPT Codes | ICD-10 Codes | HCPCS Codes | UB Revenue |
| Diabetes Screening for People with Schizophrenia or Bipolar Disorder who are using Antipsychotic Medications (SSD) (Continued from page 24) | BH Stand Alone Nonacute Inpatient Value Set with Schizophrenia, or Bipolar or Other Bipolar ICD-10 Value Set: 99304–99310, 99315, 99316, 99318, 99324–99328, 99334–99337 BH Nonacute Inpatient Value Set with BH Nonacute Inpatient POS Value Set with Schizophrenia, or Bipolar or Other Bipolar ICD-10 Value Set: 90791, 90792, 90832–90834, 90836–90840, 90845, 90847, 90849, 90853, 90867–90870, 90875, 90876, 99291 BH Nonacute Inpatient POS Value Set: 31, 32, 56 | | | |